

ON THE STRUCTURE OF THE MODULI OF JETS OF G-STRUCTURES WITH LINEAR CONNECTION.

C. MARTÍNEZ ONTALBA, J. MUÑOZ MASQUÉ, AND A. VALDÉS

ABSTRACT. The moduli space of jets of certain G -structures (basically those which admit a canonical linear connection) is shown to be isomorphic to the quotient of a natural G -module by G .

1. INTRODUCTION

Given a closed subgroup G of the general linear group $GL(n, \mathbb{R})$, a G -structure is a reduced bundle $P(M, G)$ with structure group G of the bundle of linear frames $FM \rightarrow M$.

The main types of geometries arise from different choices of G . For example, there is a one-to-one correspondence between the set of Riemannian metrics on M and the set of $O(n)$ -structures on M . Analogously, almost Hermitian geometries correspond to $U(n/2)$ -structures, almost symplectic geometries to $Sp(n/2)$ -structures, and so on.

It is a well-known fact that G -structures on M are in one-to-one correspondence with smooth sections $s \in \Gamma(FM/G)$ of the quotient bundle $FM/G \rightarrow M$. Denoting by $q : FM \rightarrow FM/G$ the natural projection, the fiber over each $x \in M$ of the G -structure $P_s \rightarrow M$ associated to $s \in \Gamma(FM/G)$ is $(P_s)_x = \{u = (x; X_1, \dots, X_n) \in F_x M \mid q(u) = s(x)\}$.

The group $\text{Diff}(M)$ of diffeomorphisms of M acts in a natural way on the set of G -structures as follows. For each diffeomorphism $f : M \rightarrow M'$, there is an associated diffeomorphism $\bar{f} : FM \rightarrow FM'$ given by $\bar{f}(x; X_1, \dots, X_n) = (f(x); f_*(X_1), \dots, f_*(X_n))$, which defines a diffeomorphism $\tilde{f} : FM/G \rightarrow FM'/G$ by $\tilde{f}[u] = [\bar{f}(u)]$. The action of $\text{Diff}(M)$ on $\Gamma(FM/G)$ is defined by:

$$f \cdot s = \tilde{f} \circ s \circ f^{-1}, \quad f \in \text{Diff}(M), \quad s \in \Gamma(FM/G).$$

Two G -structures s and s' are said to be equivalent if they are related by a diffeomorphism $f \in \text{Diff}(M)$, which amounts to the fact that $\tilde{f}(P_s) = P_{s'}$, and they are said to be locally equivalent at points $p \in M$ and $p' \in M'$ if they are equivalent in some open neighborhoods of p and p' by a diffeomorphism which maps p to p' . We call the quotient $\mathfrak{M}_G(M) = \Gamma(FM/G)/\text{Diff}(M)$ the *moduli space of G -structures* on M . The description of this space is a basic problem in differential geometry.

Date: June, 1998.

1991 Mathematics Subject Classification. Primary 53A55; Secondary 53B30.

Key words and phrases. Differential Invariant, G -structure, Moduli of G -structures.

Supported by DGICYT under grant PB95-0124.

This paper is in final form and no version of it will be submitted for publication elsewhere.

It is possible to state an analogous problem for analytic G -structures. To study this category, it is natural to introduce the spaces $J^r(FM/G)$ of jets of G -structures. The action of $\text{Diff}(M)$ on $\Gamma(FM/G)$ induces a natural action of the groupoid $J_{\text{inv}}^{r+1}(M, M)$ of $(r+1)$ -jets of diffeomorphisms of M on the space $J^r(FM/G)$ as follows:

$$(j_x^{r+1}f) \cdot (j_x^r s) = j_{f(x)}^r(f \cdot s), \quad j_x^{r+1}f \in J_{\text{inv}}^{r+1}(M, M), \quad j_x^r s \in J^r(FM/G).$$

We will call the quotient $\mathfrak{M}_G^r(M) = J^r(FM/G) / J_{\text{inv}}^{r+1}(M, M)$ the *moduli space of r -jets of G -structures* on M . There are natural projections $\mathfrak{M}_G^{r+k}(M) \rightarrow \mathfrak{M}_G^r(M)$, $k \geq 0$, so that we can define the moduli space of jets of G -structures as the projective limit $\mathfrak{M}_G^\infty(M) = \varprojlim \mathfrak{M}_G^r(M)$. The local equivalence problem for analytic G -structures can be reduced to the study of this moduli space.

Another reason for analyzing $\mathfrak{M}_G^\infty(M)$ is that it is the space where the geometric objects associated to G -structures are defined.

In general, the description of $\mathfrak{M}_G^\infty(M)$ is an extremely complicated problem. The aim of this paper is to describe, to some extent, the structure of this space in the particular case of $G \subset GL(n, \mathbb{R})$ being a closed subgroup such that the first prolongation $\mathfrak{g}^{(1)}$ vanishes and there exists a supplementary G -submodule¹ W of $\delta((\mathbb{R}^n)^* \otimes \mathfrak{g})$ in $\bigwedge^2(\mathbb{R}^n)^* \otimes \mathbb{R}^n$. As we will explain later, this is a technical condition which assures that G -structures have a canonical linear connection attached.

More precisely, we will prove the following theorem.

Theorem 1.1. *Let M be a n -dimensional smooth manifold, and let $G \subset GL(n, \mathbb{R})$ be a closed subgroup such that the first prolongation $\mathfrak{g}^{(1)}$ of its Lie algebra $\mathfrak{g}$ vanishes and there exists a supplementary G -submodule W of $\delta((\mathbb{R}^n)^* \otimes \mathfrak{g})$ in $\bigwedge^2(\mathbb{R}^n)^* \otimes \mathbb{R}^n$. Then, there exists a family of G -modules $\mathbf{S}^r$ and homomorphisms $\mathbf{S}^{r+k} \rightarrow \mathbf{S}^r$, $k \geq 0$, such that each space $\mathfrak{M}_G^r(M)$ of r -jets of G -structures is canonically isomorphic to the quotient $\mathbf{S}^r/G$.*

The moduli space $\mathfrak{M}_G^\infty(M)$ is then canonically isomorphic to the quotient $\mathbf{S}^\infty/G$, where $\mathbf{S}^\infty = \varprojlim \mathbf{S}^r$.

2. NATURAL LINEAR CONNECTIONS

In this section we will recall a classical result ([2]) which shows that, under the conditions on G stated in Theorem 1.1, it is possible to associate a linear connection to each G -structure in a natural way. We will also state certain properties of this assignment which will be useful later.

Let $G \subset GL(n, \mathbb{R})$ be a closed Lie subgroup, and $\mathfrak{g} \subset gl(n, \mathbb{R})$ its Lie algebra. Let us denote $V = \mathbb{R}^n$. Under the identification $gl(n, \mathbb{R}) \cong V^* \otimes V$, we can consider $\mathfrak{g}$ as a Lie subalgebra of the endomorphisms of V , $\text{End}(V) \cong V^* \otimes V$.

The alternation operator $\delta : V^* \otimes \mathfrak{g} \rightarrow \bigwedge^2 V^* \otimes V$ is given by

$$\delta\tau(u, v) = \tau(u)v - \tau(v)u, \quad \tau \in V^* \otimes \mathfrak{g}, u, v \in V.$$

The first prolongation of $\mathfrak{g}$ is then defined as the vector space $\mathfrak{g}^{(1)} := \ker \delta$ and, from now on, we will assume that this prolongation vanishes, i.e., $\mathfrak{g}^{(1)} = \{0\}$.

¹Throughout this paper, by G -module we mean a linear representation of G in a finite dimensional vector space.

The natural action of G on V defines linear G -actions on the spaces $V^\star \otimes \mathfrak{g}$ and $\bigwedge^2 V^\star \otimes V$, with respect to which the operator δ is an homomorphism of G -modules. The image of this operator is a G -submodule and we assume that it admits a supplementary G -submodule W :

$$(1) \quad \bigwedge^2 V^\star \otimes V = \delta(V^\star \otimes \mathfrak{g}) \oplus W.$$

Given a G -structure $s \in \Gamma(FM/G)$, the bundle $\bigwedge^2 T^\star M \otimes TM$ is identified with the associated bundle $P_s \times_G (\bigwedge^2 V^\star \otimes V)$. The decomposition (1) allows to express this bundle as the sum:

$$\bigwedge^2 T^\star M \otimes TM = (P_s \times_G \delta(V^\star \otimes \mathfrak{g})) \oplus (P_s \times_G W).$$

Let $\theta : T(FM) \rightarrow V$ be the canonical 1-form of FM , which maps each tangent vector X_u at a frame $u \in FM$ to $\theta(X_u) = u^{-1}(\pi_\star(X_u)) \in V$, where $\pi : FM \rightarrow M$ is the canonical projection and the frame u is understood as a linear isomorphism $u : V \rightarrow T_{\pi(u)}M$. We also denote by θ the restriction of the canonical form to the subbundle P_s . The restriction of the projection $\pi_\star$ to each horizontal complement $H_u \subset T_u P_s$ to the vertical tangent space at $u \in P_s$ is an isomorphism $H_u \cong T_{\pi(u)}M$, so that for any frame u and any vector $v \in V$ there is a unique $B_u(v) \in H_u$ such that $\theta(B_u(v)) = v$. The horizontal complement H_u defines then an element $t(H_u) \in \bigwedge^2 V^\star \otimes V$ by:

$$t(H_u)(v, w) = d\theta(B_u(v), B_u(w)), \quad v, w \in V.$$

If H_u are the horizontal subspaces of a linear connection on M adapted to P_s , then there is a well-defined and G -equivariant *torsion function*: $P_s \rightarrow \bigwedge^2 V^\star \otimes V$, given by $u \mapsto t(H_u)$, which is related to the torsion tensor $T \in \Gamma(\bigwedge^2 T^\star M \otimes TM)$ by: $t(H_u)(v, w) = u^{-1}(T_x(u(v), u(w)))$ for each $u \in P_s$ and $v, w \in V$.

It can be shown (see, e.g., [2] and [6]) that the condition $t(H_u) \in W$ characterizes the horizontal subspaces defining a linear connection adapted to P_s . In other words:

Theorem 2.1 ([2]). *Let $G \subset GL(n, \mathbb{R})$ be a closed Lie subgroup such that the first prolongation $\mathfrak{g}^{(1)}$ vanishes and there exists a supplementary G -submodule W of $\delta(V^\star \otimes \mathfrak{g})$ in $\bigwedge^2 V^\star \otimes V$. Then, for each G -structure $s \in \Gamma(FM/G)$ and each $u \in P_s := s(M)$ there exists a unique horizontal space $H_u \subset T_u P$ such that $t(H_u) \in W$. These horizontal spaces define a linear connection $\nabla(s)$ adapted to P_s which is characterized by the condition that its torsion is a section of the vector bundle $P_s \times_G W$.*

Next two lemmas provide some properties of this *canonical connection* which will be used to describe the structure of the moduli spaces in later sections.

Lemma 2.2. *The assignment $s \rightarrow \nabla(s)$ defines an operator $\nabla : \Gamma(FM/G) \rightarrow \Gamma(C(M))$, with $C(M) \rightarrow M$ being the affine bundle of linear connections on M , which satisfies:*

1. ∇ is natural, i.e., for each diffeomorphism $f : M \rightarrow M'$, the direct image of $\nabla(s)$ onto $\bar{f}(P_s)$ is $\nabla(f \cdot s)$.
2. $\nabla(s)$ is an operator of order 1, i.e., if $j_x^1 s = j_x^1 s'$, then $\nabla(s)(x) = \nabla(s')(x)$. Thus, ∇ defines a map, which we will also denote $\nabla : J^1(FM/G) \rightarrow C(M)$, by $\nabla(j_x^1 s) = \nabla(s)(x)$.

Proof. In order to prove the naturality of the map $\nabla : \Gamma(FM/G) \rightarrow \Gamma(C(M))$, it suffices to show that $t(H_u) \in W$ implies $t(\bar{f}_*(H_u)) \in W$ for each diffeomorphism $f : M \rightarrow M'$ because, then, the direct image $\bar{f}_*(H_u) \subset \bar{f}_*(T_u P) = T_{\bar{f}(u)} P_{f \cdot s}$ of each horizontal space of the connection $\nabla(s)$ by f is, by Theorem 2.1, a horizontal space of the connection $\nabla(f \cdot s)$. But this follows easily from the functoriality of the canonical 1-form of the bundle of linear frames. Explicitly, the canonical 1-form $\theta' : TFM' \rightarrow V$ of FM' is related to θ by $\theta = \bar{f}^* \theta'$ as one can easily check. From this it follows that $\theta'(\bar{f}_*(B_u)(v)) = \theta((B_u)(v)) = v$, for all $v \in V$, and therefore we have that $B'_{\bar{f}(u)} = \bar{f}_*(B_u)$ and so

$$\begin{aligned} t(\bar{f}_*(H_u))(v, w) &= d\theta'(B'_{\bar{f}(u)}(v), B'_{\bar{f}(u)}(w)) \\ &= d\theta'(\bar{f}_*(B_u)(v), \bar{f}_*(B_u)(w)) \\ &= d(\bar{f}^* \theta')((B_u)(v), (B_u)(w)) \\ &= d\theta((B_u)(v), (B_u)(w)) = t(H_u)(v, w). \end{aligned}$$

To prove that ∇ is an operator of order 1, first notice that the condition $j_x^1 s = j_x^1 s'$ is equivalent to the fact that the mappings $s_* : T_x M \rightarrow T_{s(x)}(FM/G)$ and $s'_* : T_x M \rightarrow T_{s'(x)}(FM/G)$ coincide. But, then, the equality $T_u P_s = T_u P_{s'}$ holds for any $u \in s(x) = s'(x)$. To see this, notice that if $q : FM \rightarrow FM/G$ is, as before, the natural projection, then:

$$\begin{aligned} T_u P_s &= q_*^{-1}(T_{q(u)}(s(M))) \\ &= q_*^{-1}(s_*(T_x M)) \\ &= q_*^{-1}(s'_*(T_x M)) \\ &= T_u P_{s'}. \end{aligned}$$

Since $t(H_u) \in W$ is an algebraic condition depending only on the vector spaces $T_u P_s$ and W , it follows that the equality $T_u P_s = T_u P_{s'}$ implies the equality of the horizontal subspaces defining the connections: $H_u = H'_u$ for $u \in s(x) = s'(x)$. ■

Now, let $\sigma : \mathcal{U} \subset M \rightarrow P_s$ be a section defined in a coordinate neighborhood $(\mathcal{U}, (x^1, \dots, x^n))$ of $x \in M$. We will denote $\sigma = (X_1, \dots, X_n)$, with $X_i = \sum_j \sigma_{ji} \frac{\partial}{\partial x^j}$ for each $i = 1, \dots, n$. Let $\omega \in \Gamma(T^* P_s \otimes \mathfrak{g})$ be the connection form of $\nabla := \nabla(s)$.

If we consider, as before, each $\sigma(x) \in P_s$ as a linear isomorphism $\sigma(x) : V \rightarrow T_x M$, we can define a function $\eta \in C^\infty(\mathcal{U}, V^* \otimes \mathfrak{g})$ by: $\eta(x) = (\sigma^* \omega)_x \circ \sigma(x)$. The components of η in the standard basis $\{v_1, \dots, v_n\}$ of V are related to those of the local connection 1-form $\sigma^* \omega \in \Gamma(T^* M \otimes \mathfrak{g})$ by: $\eta_{ij}^k = (\sigma^* \omega)_j^k(X_i)$.

Let $P_{\text{im}\delta} : \bigwedge^2 V^* \otimes V \rightarrow \delta(V^* \otimes \mathfrak{g})$ be the projection onto $\delta(V^* \otimes \mathfrak{g})$ according to the decomposition (1). Since $\ker \delta = \{0\}$, it makes sense to consider the homomorphism:

$$\delta^{-1} \circ P_{\text{im}\delta} : \bigwedge^2 V^* \otimes V \rightarrow V^* \otimes \mathfrak{g}.$$

Lemma 2.3. *The local connection 1-form $\sigma^* \omega \in \Gamma(T^* M \otimes \mathfrak{g})$ is determined by the equation: $\eta(x) = -(\delta^{-1} \circ P_{\text{im}\delta})(\tilde{t}(\sigma(x)))$, where $\tilde{t}$ is the torsion function of the linear flat connection making parallel σ .*

Proof. Using the identity $\nabla_X X_j = \sum_k X_k (\sigma^* \omega)_j^k (X)$, $X \in \Gamma(TM)$, $1 \leq j \leq n$, we see that the components of the torsion tensor of ∇ in the moving frame σ , which are given by $T(X_i, X_j) = \nabla_{X_i} X_j - \nabla_{X_j} X_i - [X_i, X_j] = \sum_k T_{ij}^k X_k$, $1 \leq i, j \leq n$, can be written as $T_{ij}^k(x) = \eta_{ij}^k(x) - \tilde{\eta}_{ji}^k(x) + \tilde{t}_{ij}^k(\sigma(x))$. Using the relation $t(H_{\sigma(x)})(v_i, v_j) = \sum_k T_{ij}^k(x) v_k$, last equation reads:

$$t(H_{\sigma(x)}) = \delta\eta(x) + \tilde{t}(\sigma(x)).$$

The condition $t(H_{\sigma(x)}) \in W$ can be written as $(\delta^{-1} \circ P_{\text{im}\delta})(t(H_{\sigma(x)})) = 0$ or, equivalently, $\eta(x) = -(\delta^{-1} \circ P_{\text{im}\delta})(\tilde{t}(\sigma(x)))$. ■

Remark 2.1. Notice that $\tilde{t}_{ij}^k(\sigma(x)) = -\sum_{h,l} \sigma^{kl} \left(\sigma_{hi} \frac{\partial \sigma_{lj}}{\partial x^h} - \sigma_{hj} \frac{\partial \sigma_{li}}{\partial x^h} \right)$, where (σ^{ij}) stands for the inverse matrix of (σ_{ij}) . Thus, the components $(\sigma^* \omega)_\beta^\gamma \in \Gamma(T^*M)$ of the local connection 1-form $\sigma^* \omega \in \Gamma(T^*M \otimes \mathfrak{g})$ are given by:

$$(2) \quad (\sigma^* \omega)_\beta^\gamma (X_\alpha) = \sum_{i,j,k,h,l} A_{\alpha\beta k}^{\gamma ij} \sigma^{kl} \left(\sigma_{hi} \frac{\partial \sigma_{lj}}{\partial x^h} - \sigma_{hj} \frac{\partial \sigma_{li}}{\partial x^h} \right), \quad 1 \leq \alpha, \beta, \gamma \leq n,$$

where the real coefficients $A_{\alpha\beta k}^{\gamma ij}$ are determined by the homomorphism $\delta^{-1} \circ P_{\text{im}\delta}$. These coefficients are then universal, in the sense that they only depend on the group $G \subset GL(n, \mathbb{R})$ and the supplementary W chosen to define the connections, but not on the particular G -structure considered.

Remark 2.2. Equation (2) shows that the map $\tilde{\nabla} : J^1(FM) \rightarrow C(M)$ defined as $\tilde{\nabla}(j_x^1 \sigma) = \nabla(q \circ \sigma)(x)$ is smooth. Since the diagram

$$\begin{array}{ccc} J^1(FM) & \xrightarrow{q^1} & J^1(FM/G) \\ \tilde{\nabla} \searrow & & \swarrow \nabla \\ & C(M) & \end{array}$$

is commutative, and the projection $q^1 : J^1(FM) \rightarrow J^1(FM/G)$ is a surjective submersion, we conclude that the map $\nabla : J^1(FM/G) \rightarrow C(M)$ is also smooth.

To end this section, we illustrate the construction of canonical connections with some examples. In all cases we will take W as the G -submodule of tensors $T \in \bigwedge^2 V^* \otimes V$ such that $\text{trace}(A \circ i_v T) = 0$ for every $A \in \mathfrak{g}$ and every $v \in V$. This is a supplementary G -submodule of the image of δ whenever $\mathfrak{g}^{(1)} = 0$ and $\mathfrak{g}$ is closed under transposition (see [10]).

Example 2.1. It is well-known that the canonical connection (in the sense of Theorem 2.1) associated to each $O(n)$ -structure or, equivalently, to each Riemannian metric is the Levi-Civita connection. That is because for $G = O(n)$ the alternation operator δ is surjective and, therefore, the supplementary submodule in equation (1) is $W = \{0\}$. The condition $\eta(x) = -\delta^{-1}(\tilde{t}(\sigma(x)))$ is equivalent to Koszul formula, as we next explain.

The operator $\delta^{-1} \circ P_{\text{im}\delta} = \delta^{-1} : \bigwedge^2 V^* \otimes V \rightarrow V^* \otimes \mathfrak{g}$ is easily obtained as follows. Let us denote by $\langle \cdot, \cdot \rangle$ the standard inner product in V , and take any $T = \delta\tau \in \bigwedge^2 V^* \otimes V$, $\tau \in V^* \otimes \mathfrak{g}$. Then, for any $u, v, w \in V$, the following relations

hold:

$$\begin{aligned}\langle T(u, v), w \rangle &= \langle \tau(u)v, w \rangle - \langle \tau(v)u, w \rangle \\ \langle T(w, u), v \rangle &= \langle \tau(w)u, v \rangle - \langle \tau(u)w, v \rangle \\ \langle T(w, v), u \rangle &= \langle \tau(w)v, u \rangle - \langle \tau(v)w, u \rangle\end{aligned}$$

Adding the three equations, taking into account that $\mathfrak{g} = \{A \in V^\star \otimes V \mid \langle Au, v \rangle + \langle Av, u \rangle = 0 \ \forall u, v \in V\}$, we obtain the expression of $\tau = \delta^{-1}T$:

$$(3) \quad \langle \tau(u)v, w \rangle = \frac{1}{2} (\langle T(u, v), w \rangle + \langle T(w, u), v \rangle + \langle T(w, v), u \rangle),$$

for all $u, v, w \in V$. Hence, the condition $\eta(x) = -\delta^{-1}(\tilde{t}(\sigma(x)))$ can be written, in components, as:

$$(4) \quad \eta_{ij}^k(x) = -\frac{1}{2} (\tilde{t}_{ij}^k(\sigma(x)) + \tilde{t}_{ki}^j(\sigma(x)) + \tilde{t}_{kj}^i(\sigma(x))).$$

Now, let (M, g) be a Riemannian manifold and $P_s \rightarrow M$ the corresponding $O(n)$ -structure, that is, the subbundle of orthonormal frames with respect to g . For any section $\sigma = (X_1, \dots, X_n) : \mathcal{U} \rightarrow P_s$, the components of $\eta \in C^\infty(\mathcal{U}, V^\star \otimes \mathfrak{g})$ are given by: $\eta_{ij}^k = (\sigma^\star \omega)_j^k(X_i) = g(\nabla_{X_i} X_j, X_k)$, whereas those of $\tilde{t}(\sigma(x))$ are: $\tilde{t}_{ij}^k = -g([X_i, X_j], X_k)$. Substitution of these expressions in (4) yields:

$$2g(\nabla_{X_i} X_j, X_k) = g([X_i, X_j], X_k) + g([X_k, X_i], X_j) + g([X_k, X_j], X_i),$$

which is nothing but Koszul formula in the orthonormal frame $\{X_1, \dots, X_n\}$.

Example 2.2. Consider now the case of $(O(p) \times O(q))$ -structures, i.e., those with structure group $G = \left\{ \begin{pmatrix} A_1 & 0 \\ 0 & A_2 \end{pmatrix} \in O(n) \mid A_1 \in O(p), A_2 \in O(q) \right\}$. These structures correspond to Riemannian almost product structures.

Let $\{v_1, \dots, v_n\}$ be the canonical basis of V , and let us denote $I_1 = \{1, \dots, p\}$, $I_2 = \{p+1, \dots, p+q\}$, $V_\alpha = \text{Span}\{v_i\}_{i \in I_\alpha}$, $\alpha = 1, 2$. It can be easily seen that W is given as the G -submodule of tensors $\tilde{T} \in \bigwedge^2 V^\star \otimes V$ such that:

$$(5) \quad \langle \tilde{T}(u, v), w \rangle = \langle \tilde{T}(u, w), v \rangle \text{ if } v, w \in V_\alpha, \ \alpha = 1, 2,$$

for all $u \in V$.

In order to determine an expression of the operator $\delta^{-1} \circ P_{\text{im} \delta}$, let us decompose any $T \in \bigwedge^2 V^\star \otimes V$ as $T = \tilde{T} + \delta \tau$, with $\tilde{T} \in W$ and $\tau \in V^\star \otimes \mathfrak{g}$.

If $v \in V_\alpha$, $w \in V_\beta$, with $\alpha \neq \beta$, then $\langle \tau(u)v, w \rangle = 0$ for each $u \in V$, because $\tau(u)$ belongs to $\mathfrak{g} = o(p) \oplus o(q)$.

If $u, v, w \in V_\alpha$, $\alpha = 1, 2$, then $\langle \tilde{T}(u, v), w \rangle = 0$, as follows easily from (5). In this case, one can compute, as in the previous example:

$$\langle \tau(u)v, w \rangle = \frac{1}{2} (\langle T(u, v), w \rangle + \langle T(w, u), v \rangle + \langle T(w, v), u \rangle).$$

Finally, if $u \in V_\alpha$ and $v, w \in V_\beta$, $\alpha \neq \beta$, we have:

$$\begin{aligned}\langle T(u, v), w \rangle &= \langle \tau(u)v, w \rangle + \langle \tilde{T}(u, v), w \rangle \\ \langle T(u, w), v \rangle &= \langle \tau(u)w, v \rangle + \langle \tilde{T}(u, w), v \rangle.\end{aligned}$$

Subtracting the second equation from the first one, we obtain:

$$\langle \tau(u)v, w \rangle = \frac{1}{2} (\langle T(u, v), w \rangle - \langle T(u, w), u \rangle).$$

Thus, in components, we have:

$$((\delta^{-1} \circ P_{\text{im}\delta})(T))_{ij}^k = \begin{cases} \frac{1}{2} (T_{ij}^k + T_{ki}^j + T_{kj}^i) & \text{if } i, j, k \in I_\alpha, \alpha = 1, 2, \\ \frac{1}{2} (T_{ij}^k - T_{ik}^j) & \text{if } i \in I_\alpha, j, k \in I_\beta, \alpha \neq \beta, \\ 0 & \text{if } j \in I_\alpha, k \in I_\beta, \alpha \neq \beta. \end{cases}$$

Let us consider now a Riemannian almost product manifold, i.e., a Riemannian manifold (M, g) and a tensor field $\phi \in \mathfrak{T}_1^1(M)$ such that $\phi^2 = \text{id}$ and $g(\phi X, \phi Y) = g(X, Y)$ for any vector fields X, Y on M . The tensor ϕ gives rise to two mutually orthogonal distributions V_1 and V_2 , corresponding to its eigenvalues 1 and -1, called the vertical and horizontal distributions, respectively. Denoting $p = \dim V_1$, $q = \dim V_2$, the corresponding $(O(p) \times O(q))$ -structure is the subbundle of orthonormal frames $(X_1, \dots, X_n)$ such that $X_1, \dots, X_p$ are vertical vectors and $X_{p+1}, \dots, X_n$ are horizontal vectors.

From the expression of $\delta^{-1} \circ P_{\text{im}\delta}$, we obtain, as in the preceding example:

$$g(\nabla_{X_i} X_j, X_k) = \frac{1}{2} (g([X_i, X_j], X_k) + g([X_k, X_i], X_j) + g([X_k, X_j], X_i))$$

if $i, j, k \in I_\alpha, \alpha = 1, 2$,

$$g(\nabla_{X_i} X_j, X_k) = \frac{1}{2} (g([X_i, X_j], X_k) - g([X_i, X_k], X_j))$$

if $i \in I_\alpha, j, k \in I_\beta, \alpha \neq \beta$, and

$$g(\nabla_{X_i} X_j, X_k) = 0$$

if $j \in I_\alpha, k \in I_\beta, \alpha \neq \beta$.

The third equation implies that ∇_X leaves the vertical and horizontal distributions invariant for all X . The first one gives:

$$g(\nabla_X Y, Z) = \frac{1}{2} (X(g(Y, Z)) + Y(g(Z, X)) - Z(g(X, Y)) + g([X, Y], Z) + g([Z, X], Y) - g([Y, Z], X)),$$

if $X, Y, Z \in V_\alpha, \alpha = 1, 2$. Denoting by ∇^{LC} the Levi-Civita connection, this means that, if $X, Y \in V_\alpha$, then $\nabla_X Y$ is the orthogonal projection of $\nabla_X^{\text{LC}} Y$ onto V_α according to the decomposition $TM = V_1 \oplus V_2$. Finally, the second equation yields:

$$g(\nabla_X Y, Z) = \frac{1}{2} (X(g(Y, Z)) + g([X, Y], Z) - g([X, Z], Y)),$$

if $X \in V_\alpha, Y, Z \in V_\beta, \alpha \neq \beta$.

Example 2.3. Next, we consider the case $G = \mathbb{R}^* \cdot I_n$ with $n \geq 2$. If $n = 2$ we obtain classical webs over surfaces (see [3]).

The Lie algebra $\mathfrak{g}$ of G is generated by the identity matrix I_n , hence we can take:

$$W = \{T \in \wedge^2 V^* \otimes V \mid \text{trace}(i_v T) = 0 \forall v \in V\}.$$

A short computation shows that the operator $\delta^{-1} \circ P_{\text{im}\delta}$ is defined by:

$$((\delta^{-1} \circ P_{\text{im}\delta})(T))(v) = \left(\frac{1}{n-1} \text{trace}(i_v T) \right) I_n,$$

for all $T \in \bigwedge^2 V^* \otimes V$, $v \in V$. The condition $\eta(x) = -\delta^{-1}(\tilde{t}(\sigma(x)))$ is written, in components:

$$\eta_{ij}^k(x) = -\frac{\delta_j^k}{n-1} \sum_l \tilde{t}_{il}^l(\sigma(x)).$$

From this expression, it follows that the canonical connection is given, in any local section $\sigma = (X_1, \dots, X_n)$, by:

$$\nabla_{X_i} X_j = \left(\frac{1}{n-1} \sum_k C_{ik}^k \right) X_j,$$

with the coefficients C_{ij}^k being defined by $[X_i, X_j] = \sum_k C_{ij}^k X_k$.

3. CANONICAL REPRESENTATION OF THE MODULI SPACES

In this section we are going to describe the underlying manifolds of the G -modules $\mathbf{S}^r$ to which we refer in Theorem 1.1, and the action of the group G on them. However, the definition of linear structures on these G -manifolds will be postponed until section 5. We also describe the isomorphisms $\mathfrak{M}_G^r(M) \cong \mathbf{S}^r/G$.

Let us denote by $\mathfrak{G}_0^{r+1} \subset J_{\text{inv}}^{r+1}(V, V)$ the Lie group of $(r+1)$ -jets of diffeomorphisms of V which leave the origin $0 \in V$ fixed. The restriction to $\mathfrak{G}_0^{r+1}$ of the action of the Lie groupoid $J_{\text{inv}}^{r+1}(V, V)$ on $J^r(FV/G)$ defines an action of $\mathfrak{G}_0^{r+1}$ on $J_0^r(FV/G)$.

Proposition 3.1. *There exists a canonical bijection $\mathfrak{M}_G^r(M) \cong J_0^r(FV/G)/\mathfrak{G}_0^{r+1}$.*

Proof. Let us recall that $\mathfrak{M}_G^r(M)$ has been defined as the quotient

$$J^r(FM/G)/J_{\text{inv}}^{r+1}(M, M).$$

For each element $[j_x^r s] \in \mathfrak{M}_G^r(M)$, we choose a chart $\varphi : \mathcal{U} \subset M \rightarrow V$, centered at $x \in M$, and define the element $(j_x^{r+1} \varphi) \cdot (j_x^r s) = j_0^r(\tilde{\varphi} \circ s \circ \varphi^{-1}) \in J_0^r(FV/G)$. If we take another representative $(j_x^{r+1} f) \cdot (j_x^r s)$ of $[j_x^r s]$, and a chart $(\mathcal{U}', \varphi')$ centered at $f(x) \in M$, we have:

$$(j_{f(x)}^{r+1} \varphi') \cdot ((j_x^{r+1} f) \cdot (j_x^r s)) = (j_0^{r+1}(\varphi' \circ f \circ \varphi^{-1})) \cdot ((j_x^{r+1} \varphi) \cdot (j_x^r s)).$$

Thus, the element of $J_0^r(FV/G)$ assigned to $[j_x^r s]$ is determined up to an $(r+1)$ -jet of the form $j_0^{r+1}(\varphi' \circ f \circ \varphi^{-1})$, which is an element of $\mathfrak{G}_0^{r+1}$ and, hence, there is a well defined mapping $\mathfrak{M}_G^r(M) \rightarrow J_0^r(FV/G)/\mathfrak{G}_0^{r+1}$. It is easy to find the inverse of this mapping. Given a class $[j_0^r t] \in J_0^r(FV/G)/\mathfrak{G}_0^{r+1}$ and a chart $\varphi : \mathcal{U} \subset M \rightarrow V$ centered at $x \in M$ the element $[(j_0^{r+1} \varphi^{-1}) \cdot (j_0^r t)] \in J^r(FM/G)/J_{\text{inv}}^{r+1}(M, M)$ is well-defined. The mapping $[j_0^r t] \mapsto [(j_0^{r+1} \varphi^{-1}) \cdot (j_0^r t)]$ is the required inverse $J_0^r(FV/G)/\mathfrak{G}_0^{r+1} \rightarrow \mathfrak{M}_G^r(M)$. ■

From now on, we will denote the spaces $\mathfrak{M}_G^r(M)$ just by $\mathfrak{M}_G^r$, because we have seen that they actually do not depend on the base manifold M .

Now, we define the spaces $\mathcal{E}^r(V)$ of *framed r -jets of G -structures* as:

$$\mathcal{E}^r(V) := J_0^r(FV/G) \times_{F_0 V/G} F_0 V = \{(j_0^r s, u) \in J_0^r(FV/G) \times F_0 V : [u] = s(0)\}.$$

We consider the left action of $\mathfrak{G}_0^{r+1}$ on $\mathcal{E}^r(V)$ given by:

$$(j_0^{r+1} f) \cdot (j_0^r s, u) = ((j_0^{r+1} f) \cdot (j_0^r s), \bar{f}(u)), \quad j_0^{r+1} f \in \mathfrak{G}_0^{r+1}, (j_0^r s, u) \in \mathcal{E}^r(V).$$

Besides, we consider the right action of G on $\mathcal{E}^r(V)$ induced by the action of G on F_0V , i.e.,

$$(j_0^r s, u) \cdot g = (j_0^r s, u \cdot g), \quad g \in G, (j_0^r s, u) \in \mathcal{E}^r(V).$$

The quotient $\mathcal{E}^r(V)/G$ can obviously be identified with $J_0^r(FV/G)$.

It is easy to check that the group actions just defined commute. Therefore, we have the bijections:

$$\mathfrak{M}_G^r \cong \frac{J_0^r(FV/G)}{\mathfrak{G}_0^{r+1}} \cong \frac{\mathcal{E}^r(V)/G}{\mathfrak{G}_0^{r+1}} \cong \frac{\mathcal{E}^r(V)/\mathfrak{G}_0^{r+1}}{G}.$$

Thus, if there is a manifold $\mathbf{S}^r$ such that $\mathbf{S}^r/G \cong \mathfrak{M}_G^r$, the natural candidate is $\mathbf{S}^r := \mathcal{E}^r(V)/\mathfrak{G}_0^{r+1}$, if it is indeed a smooth manifold without singularities. In general, this would not be true. However, the existence of linear connections functorially attached to G -structures makes each $\mathcal{E}^r(V)$ a trivial principal bundle with structure group $\mathfrak{G}_0^{r+1}$, as we will see below, and so $\mathcal{E}^r(V)/\mathfrak{G}_0^{r+1}$ is a manifold in our particular cases.

Given $(j_0^r s, u) \in \mathcal{E}^r(V)$, let $s : \mathcal{U} \subset V \rightarrow FV/G$ be a representative of $j_0^r s$, defined on an open neighborhood $\mathcal{U}$ of $0 \in V$. The connection $\nabla(s)$ provides an exponential mapping $\exp_s : \mathcal{W}_0 \subset T_0V \rightarrow V$ defined on some neighborhood $\mathcal{W}_0$ of $0 \in T_0V$. The composition of this mapping with the isomorphism $u : V \rightarrow T_0V$ yields a diffeomorphism defined on a neighborhood of $0 \in V$, which is nothing but the set of normal coordinates associated to the connection $\nabla(s)$ and the frame u .

Lemma 3.2. *The assignment $\text{Exp}^r : \mathcal{E}^r(V) \rightarrow \mathfrak{G}_0^{r+1}$ given by $\text{Exp}^r(j_0^r s, u) = j_0^{r+1}(\exp_s \circ u)$ is a well-defined smooth map. Moreover, it is $\mathfrak{G}_0^{r+1}$ -equivariant with respect to the action of the group on $\mathcal{E}^r(V)$ defined above and the natural left action of $\mathfrak{G}_0^{r+1}$ on itself.*

Proof. Let $(V, (x^1, \dots, x^n))$ be the standard chart of V . Let us denote $f = \exp_s \circ u$, and $u = (X_1, \dots, X_n)$, with $X_i = \sum_j a_i^j \left(\frac{\partial}{\partial x^j}\right)_0$, $i = 1, \dots, n$.

The map $t \mapsto f(tx)$ is the geodesic from $0 \in V$ with initial speed $u(x)$, i.e., the solution of the system of second order differential equations:

$$(6) \quad \frac{d^2 f^k(tx)}{dt^2} = - \sum_{j_1, j_2} \Gamma_{j_1 j_2}^k(f(tx)) \frac{df^{j_1}(tx)}{dt} \frac{df^{j_2}(tx)}{dt}, \quad 1 \leq k \leq n,$$

with initial conditions $f(0) = 0$, $\left.\frac{df(tx)}{dt}\right|_0 = u(x)$. We can write this system in an equivalent way as follows:

$$\begin{aligned} f(0) &= 0 \\ \sum_i x^i \frac{\partial f^k}{\partial x^i}(0) &= \sum_i x^i a_i^k, \quad 1 \leq k \leq n, \\ \sum_{i_1, i_2} x^{i_1} x^{i_2} \frac{\partial^2 f^k}{\partial x^{i_1} \partial x^{i_2}}(tx) &= - \sum_{i_1, i_2, j_1, j_2} x^{i_1} x^{i_2} \Gamma_{j_1 j_2}^k(f(tx)) \frac{\partial f^{j_1}}{\partial x^{i_1}}(tx) \frac{\partial f^{j_2}}{\partial x^{i_2}}(tx), \quad 1 \leq k \leq n. \end{aligned}$$

From the second equation it follows that $\frac{\partial f^k}{\partial x^i}(0) = a_i^k$, $1 \leq i, k \leq n$. On the other hand, taking the $(r-1)$ th order derivative ($r \geq 1$) of the second equation and

evaluating in $t = 0$ leads to the identity:

$$\sum_{i_1, \dots, i_{r+1}} x^{i_1} \dots x^{i_{r+1}} \frac{\partial^{r+1} f^k}{\partial x^{i_1} \dots \partial x^{i_{r+1}}}(0) = \sum_{i_1, \dots, i_{r+1}} x^{i_1} \dots x^{i_{r+1}} R_{i_1 \dots i_{r+1}}, \quad 1 \leq k \leq n,$$

where each $R_{i_1 \dots i_{r+1}}$ is a polynomial in the derivatives of the Christoffel symbols, at $x = 0$, up to order $r - 1$, and the derivatives at $x = 0$ of the components f^k up to order r .

Last equation implies:

$$\frac{\partial^{r+1} f^k}{\partial x^{i_1} \dots \partial x^{i_{r+1}}}(0) = \frac{1}{(r+1)!} \sum_{\tau \in S_{r+1}} R_{i_{\tau(1)} \dots i_{\tau(r+1)}}, \quad 1 \leq k, i_1, \dots, i_{r+1} \leq n.$$

Since the map $j_x^1 s \mapsto \nabla(s)(x)$ is smooth (see Remark 2.2), the derivatives up to order $r - 1$ of the Christoffel symbols $\Gamma_{ij}^k(x)$ are smooth functions of $j_x^r s$. Thus, the $(r+1)$ th order derivatives at $x = 0$ of f are smooth functions of $j_0^r s$ and $j_0^r f$.

Finally, using induction, it follows that $\text{Exp}^r : (j_0^r s, u) \mapsto j_0^{r+1} f$ is a well-defined smooth map.

In order to prove the $\mathfrak{G}_0^{r+1}$ -equivariance, let us point that, due to the naturality of the assignment $s \mapsto \nabla(s)$, the equality $f \circ \exp_s \circ u = \exp_{f \cdot s} \circ \bar{f}(u)$ holds for each diffeomorphism $f : M \rightarrow M'$ and each G -structure $s \in \Gamma(FM/G)$. Therefore:

$$\begin{aligned} \text{Exp}^r((j_0^{r+1} f) \cdot (j_0^r s, u)) &= \text{Exp}^r(j_0^r(f \cdot s), \bar{f}(u)) \\ &= j_0^{r+1}(\exp_{f \cdot s} \circ \bar{f}(u)) \\ &= j_0^{r+1}(f \circ \exp_s \circ u) \\ &= (j_0^{r+1} f) \cdot \text{Exp}^r(j_0^r s, u), \end{aligned}$$

for each $j_0^{r+1} f \in \mathfrak{G}_0^{r+1}$ and each $(j_0^r s, u) \in \mathcal{E}^r(V)$. ■

Next, we define $\mathcal{E}^r(V)_1$ as the fiber over $1_{\mathfrak{G}_0^{r+1}} \in \mathfrak{G}_0^{r+1}$ of the map $\text{Exp}^r : \mathcal{E}^r(V) \rightarrow \mathfrak{G}_0^{r+1}$, i.e., $\mathcal{E}^r(V)_1 := (\text{Exp}^r)^{-1}(1_{\mathfrak{G}_0^{r+1}})$.

Lemma 3.3. $\mathcal{E}^r(V)_1$ is a smooth submanifold of $\mathcal{E}^r(V)$.

Proof. We prove first that, due to equivariance, the map Exp^r is a surjective submersion, which implies that $\mathcal{E}^r(V)_1$ is in fact a submanifold of $\mathcal{E}^r(V)$. It is surjective because the image of each orbit in $\mathcal{E}^r(V)$ is an orbit of left translations in the group, namely, $\mathfrak{G}_0^{r+1}$. To see that it is a submersion, note that if $(j_0^r s, u) \in \mathcal{E}^r(V)$ were a critical point of Exp^r , then every element of its orbit would also be a critical point, and, being surjective restricted to each orbit, the map Exp^r would not have any regular value, contradicting Sard's Theorem. Thus, each fiber of Exp^r , in particular $\mathcal{E}^r(V)_1$, is a smooth submanifold of $\mathcal{E}^r(V)$. ■

On the other hand, the equivariance of Exp^r allows to define a map $p^r : \mathcal{E}^r(V) \rightarrow \mathcal{E}^r(V)_1$ as

$$p^r(j_0^r s, u) = (\text{Exp}^r(j_0^r s, u))^{-1} \cdot (j_0^r s, u).$$

A straightforward calculation shows that this map is invariant under the right action associated to the left action of $\mathfrak{G}_0^{r+1}$ on $\mathcal{E}^r(V)$, which is given by:

$$(j_0^r s, u) \cdot (j_0^{r+1} f) = (j_0^{r+1} f)^{-1} \cdot (j_0^r s, u), \quad j_0^{r+1} f \in \mathfrak{G}_0^{r+1}, (j_0^r s, u) \in \mathcal{E}^r(V).$$

Indeed, the following lemma holds:

Lemma 3.4. $p^r : \mathcal{E}^r(V) \rightarrow \mathcal{E}^r(V)_1$ is a trivial principal bundle with structure group $\mathfrak{G}_0^{r+1}$.

Proof. The map $\Lambda^r : \mathcal{E}^r(V) \rightarrow \mathcal{E}^r(V)_1 \times \mathfrak{G}_0^{r+1}$ defined by: $\Lambda^r(j_0^r s, u) = (p^r(j_0^r s, u), (\text{Exp}^r(j_0^r s, u))^{-1})$ is a $\mathfrak{G}_0^{r+1}$ -equivariant diffeomorphism. The inverse $(\Lambda^r)^{-1} : \mathcal{E}^r(V)_1 \times \mathfrak{G}_0^{r+1} \rightarrow \mathcal{E}^r(V)$ is given by $(\Lambda^r)^{-1}((j_0^r s, u), j_0^{r+1} f) = (j_0^r s, u) \cdot (j_0^{r+1} f)$. This diffeomorphism endows $\mathcal{E}^r(V)$ with the structure of a trivial principal $\mathfrak{G}_0^{r+1}$ -bundle over $\mathcal{E}^r(V)_1$. ■

Corollary 3.5. The space $\mathbf{S}^r := \mathcal{E}^r(V)/\mathfrak{G}_0^{r+1}$ is a smooth manifold canonically diffeomorphic to $\mathcal{E}^r(V)_1$.

Remark 3.1. Explicitly, the diffeomorphism $\bar{p}^r : \mathbf{S}^r \rightarrow \mathcal{E}^r(V)_1$ is given by

$$\bar{p}^r([(j_0^r s, u)]) = p^r(j_0^r s, u).$$

Its inverse is the composed map: $\mathcal{E}^r(V)_1 \hookrightarrow \mathcal{E}^r(V) \rightarrow \mathcal{E}^r(V)/\mathfrak{G}_0^{r+1}$ of the inclusion (which is the map trivializing the bundle) with the natural projection.

Remark 3.2. Let us denote by $u^0 : V \rightarrow T_0 V$ the canonical frame. The condition $j_0^{r+1}(\exp_s \circ u) = 1_{\mathfrak{G}_0^{r+1}}$ defining $\mathcal{E}^r(V)_1$ implies $D(\exp_s \circ u)(0) = \text{id}_V$, where D stands for the standard Fréchet derivative in V . Using that

$$\begin{aligned} D(\exp_s \circ u)(0) &= D(\exp_s \circ u^0)(0) \circ D((u^0)^{-1} \circ u)(0) = \\ &= D((u^0)^{-1} \circ u)(0) = (u^0)^{-1} \circ u, \end{aligned}$$

we conclude that $u = u^0$, so that $\mathcal{E}^r(V)_1 \subset J_0^r(FV/G) \times \{u^0\}$. Therefore from now on, we will consider $\mathcal{E}^r(V)_1$ as a subspace of $J_0^r(FV/G)$.

Remark 3.3. Notice that an r -jet $j_0^r s \in J_0^r(FV/G)$ belongs to $\mathcal{E}^r(V)_1$ if and only if $s(0) = [u^0]$ and

$$(7) \quad \left(\frac{d^{k-1}}{dt^{k-1}} \sum_{i_1, i_2} x^{i_1} x^{i_2} \Gamma_{i_1 i_2}^\gamma(tx) \right) (0) = 0, \quad 1 \leq k \leq r, \quad 1 \leq \gamma \leq n,$$

for any x in a neighborhood of $0 \in V$, as follows from (6).

The action of G on $\mathbf{S}^r$ can be translated now to $\mathcal{E}^r(V)_1$.

Lemma 3.6. The action of G on $\mathbf{S}^r$ induces, by means of the diffeomorphism $\mathbf{S}^r \cong \mathcal{E}^r(V)_1$, the following action of G on $\mathcal{E}^r(V)_1$:

$$(j_0^r s) \cdot g = (j_0^{r+1} g^{-1}) \cdot j_0^r s, \quad j_0^r s \in \mathcal{E}^r(V)_1, \quad g \in G \subset \mathfrak{G}_0^{r+1}.$$

Proof. For each $j_0^r s \in \mathcal{E}^r(V)_1$ and each $g \in G$, we have, taking into account Remark 3.1 we have that

$$\begin{aligned} (j_0^r s) \cdot g &\doteq \bar{p}^r([(j_0^r s, u^0)] \cdot g) \\ &= \bar{p}^r([(j_0^r s, u^0 \circ g)]) \\ &= p^r(j_0^r s, u^0 \circ g) \\ &= (\text{Exp}^r(j_0^r s, u^0 \circ g))^{-1} \cdot (j_0^r s, u^0 \circ g). \end{aligned}$$

From the fact that:

$$\begin{aligned} \text{Exp}^r(j_0^r s, u \circ g) &= j_0^{r+1}(\exp_s \circ u^0 \circ g) \\ &= j_0^{r+1}(\exp_s \circ u^0) \cdot j_0^{r+1}g \\ &= j_0^{r+1}g, \end{aligned}$$

it follows the desired expression of $(j_0^r s) \cdot g$:

$$\begin{aligned} (j_0^r s) \cdot g &= (j_0^{r+1}g^{-1}) \cdot (j_0^r s, u^0 \circ g) \\ &= ((j_0^{r+1}g^{-1}) \cdot (j_0^r s), u^0) \end{aligned}$$

which is identified with $(j_0^{r+1}g^{-1}) \cdot (j_0^r s)$. ■

It is immediate that the projections $J_0^{r+k}(FV/G) \rightarrow J_0^r(FV/G)$ induce projections $\mathbf{S}^{r+k} \rightarrow \mathbf{S}^r$, $k \geq 0$, which allow to define $\mathbf{S}^\infty = \varprojlim \mathbf{S}^r$ as the projective limit of the family $\{\mathbf{S}^r\}$. The action of G induces an action on the limit $\mathbf{S}^\infty$ in a natural way.

Thus, we have proved the following theorem of reduction of the group:

Theorem 3.7. *The spaces $\mathbf{S}^r = \mathcal{E}^r(V)/\mathfrak{G}_0^{r+1}$ have natural smooth G -manifold structures, so that each moduli space $\mathfrak{M}_G^r$ is canonically isomorphic to the quotient $\mathbf{S}^r/G$. Taking projective limits yields a canonical isomorphism $\mathfrak{M}_G^\infty \cong \mathbf{S}^\infty/G$.*

4. EMBEDDING OF THE MODULI OF FRAMED r -JETS $\mathbf{S}^r$ IN THE SPACE OF JETS OF MOVING FRAMES $J_0^r(FV)$

Let us denote by $J_0^r(FV/G)_{u^0}$ the manifold of r -jets $j_0^r s \in J_0^r(FV/G)$ such that $s(0) = [u^0]$. We will define an embedding of this manifold in $J_0^r(FV)$. Then, from the diffeomorphism $\mathbf{S}^r \cong \mathcal{E}^r(V)_1$ and the inclusion $\mathcal{E}^r(V)_1 \hookrightarrow J_0^r(FV/G)_{u^0}$, we will obtain an embedding of $\mathbf{S}^r$ in $J_0^r(FV)$. This construction is a technical tool to define the G -module structure on $\mathbf{S}^r$ in the next section.

We define $\Upsilon^r : J_0^r(FV/G)_{u^0} \rightarrow J_0^r(FV)$ as follows. Given $j_0^r s \in J_0^r(FV/G)_{u^0}$, let $s : \mathcal{U} \subset V \rightarrow FV/G$ be a local representative of $j_0^r s$. We define a moving frame $\sigma = (X_1, \dots, X_n)$ in a neighborhood of $0 \in V$ as the $\nabla(s)$ -parallel transport of the frame u^0 along each ray $x(t) = (t \cdot \lambda^1, \dots, t \cdot \lambda^n) \in V$, and we set $\Upsilon^r(j_0^r s) = j_0^r \sigma$.

Lemma 4.1. *The assignment $\Upsilon^r : J_0^r(FV/G)_{u^0} \rightarrow J_0^r(FV)$ is a well-defined embedding.*

Proof. Let s be a representative of $j_0^r s$ defined in a neighborhood of $0 \in V$, and denote by Γ_{ij}^k the Christoffel symbols of the connection $\nabla(s)$. Let $\sigma = (X_1, \dots, X_n)$ be the moving frame defined above using the connection $\nabla(s)$, and let $X_i = \sum_k \sigma_{ki}(x) \partial/\partial x^k$ be the expression of the vector field X_i with respect to the standard chart of V .

Since, by definition, each X_i is parallel along the rays $x(t) = t\lambda$, $\lambda = (\lambda^1, \dots, \lambda^n)$, the functions σ_{ki} satisfy the differential equations:

$$\begin{aligned} \sum_{\beta} \left(\frac{\partial \sigma_{\gamma i}}{\partial x^{\beta}}(t\lambda) + \sum_{\alpha} \Gamma_{\beta\alpha}^{\gamma}(t\lambda) \sigma_{\alpha i}(t\lambda) \right) \lambda^{\beta} &= 0 \\ \sigma_{ij}(0) &= \delta_{ij} \end{aligned}$$

for any $\lambda \in V$, and any indices $1 \leq \gamma, i, j \leq n$. Multiplying these equations by t , we obtain that σ is characterized by the initial condition $\sigma(0) = u^0$ and the equations:

$$(8) \quad \sum_{\beta} \left(\frac{\partial \sigma_{\gamma i}}{\partial x^{\beta}} + \sum_{\alpha} \Gamma_{\beta \alpha}^{\gamma} \sigma_{\alpha i} \right) x^{\beta} = 0, \quad 1 \leq i, \gamma \leq n.$$

Iterated derivation of (8) and evaluation in $x = 0$ leads to:

$$(9) \quad \frac{\partial^k \sigma_{\gamma i}}{\partial x^{i_1} \dots \partial x^{i_k}}(0) = -\frac{1}{k_{(i_1 \dots i_k)}} \mathfrak{S} \left(\frac{\partial^{k-1}}{\partial x^{i_2} \dots \partial x^{i_k}} \sum_{\alpha} \Gamma_{i_1 \alpha}^{\gamma} \sigma_{\alpha i} \right)(0),$$

for any $k \geq 1$ and any $1 \leq \gamma, i, i_1, \dots, i_r \leq n$, where $\mathfrak{S}$ stands for the cyclic sum with respect to the corresponding indices.

For $r = 1$ we have $\frac{\partial \sigma_{\gamma i}}{\partial x^j}(0) = -\Gamma_{ji}^{\gamma}(0)$, so that $j_0^1 \sigma = \Upsilon^1(j_0^1 s)$ depends smoothly on $j_0^1 s$. By induction, using (9), it follows that $j_0^r \sigma = \Upsilon^r(j_0^r s)$ depends smoothly on $j_0^r s$.

Finally, it is obvious, by its very definition, that Υ^r takes its values in the closed submanifold: $J_0^r(FV)_{u^0} = \{j_0^r \sigma \in J_0^r(FV) \mid \sigma(0) = u^0\}$, and that it is a section of the projection $J_0^r(FV)_{u^0} \rightarrow J_0^r(FV/G)_{u^0}$ induced by $q : FV \rightarrow FV/G$. Therefore, Υ^r is indeed an embedding. ■

Next we define a right action of G on $J_0^r(FV)_{u^0}$ by:

$$(j_0^r \sigma) \cdot g = j_0^r (R_g \circ (g^{-1} \cdot \sigma)), \quad j_0^r \sigma \in J_0^r(FV)_{u^0}, \quad g \in G,$$

where R_g stands for right translations in FV . Besides, we have the right action of G on $J_0^r(FV/G)_{u^0}$:

$$(j_0^r s) \cdot g = (j_0^{r+1} g^{-1}) \cdot j_0^r s, \quad j_0^r s \in J_0^r(FV/G)_{u^0}, \quad g \in G.$$

Lemma 4.2. *The map $\Upsilon^r : J_0^r(FV/G)_{u^0} \rightarrow J_0^r(FV)_{u^0}$ is G -equivariant with respect to the right actions of G previously defined.*

Proof. Due to the naturality of the assignment $s \mapsto \nabla(s)$, if σ is the parallel moving frame associated to the G -structure s , then $R_g \circ (g^{-1} \cdot \sigma)$ is the parallel moving frame associated to $s \cdot g = \tilde{g}^{-1} \circ s \circ g$. From this fact, the result follows easily. ■

Denoting $\mathcal{S}^r = \Upsilon^r(\mathcal{E}^r(V)_1) \subset J_0^r(FV)$ we have now the identifications between G -manifolds $\mathbf{S}^r \cong \mathcal{E}^r(V)_1 \cong \mathcal{S}^r$.

We derive now a characterization of the submanifolds $\mathcal{S}^r \subset J_0^r(FV)$, which will be used in next section to define inductively a G -module structure on each of them. First of all, we introduce some definitions.

Let us denote $V_q^{r,p} := \mathcal{S}^r(V^*) \otimes V^{\star \otimes p} \otimes V^{\otimes q}$ with its natural G -module structure. Consider the homomorphisms of G -modules $\text{sym}_q^{r,p} : V_q^{r,p} \rightarrow V_q^{r+1,p-1}$, $p \geq 1$, given by symmetrization on the first $r+1$ covariant indices. We will denote $\bar{V}_q^{r,p} := \ker(\text{sym}_q^{r,p})$.

Let $\sigma = (X_1, \dots, X_n)$, with $X_i = \sum_j \sigma_{ji} \partial / \partial x^j$, $1 \leq i \leq n$, be a representative of the r -jet $j_0^r \sigma \in J_0^r(FV)$. Then, for each $g \in G$, the derivatives of the components of the vectors in the representative $R_g \circ (g^{-1} \cdot \sigma)$ of $(j_0^r \sigma) \cdot g$ are given by

$$\frac{\partial^k \bar{\sigma}_{rs}}{\partial x^{p_1} \dots \partial x^{p_k}}(0) = g^{r\alpha} \frac{\partial^k \sigma_{\alpha\gamma}}{\partial x^{l_1} \dots \partial x^{l_k}}(0) g_{l_1 p_1} \dots g_{l_k p_k} g_{\gamma s}.$$

Thus, we can define an equivariant mapping $\Sigma^{(r)} : J_0^r(FV) \rightarrow V_1^{r,1}$ by:

$$\Sigma^{(r)}(j_0^r \sigma) = \sum_{l_1, \dots, l_r} \sum_{i,j} \frac{\partial^r \sigma_{ij}}{\partial x^{l_1} \dots \partial x^{l_r}}(0) v^{*l_1} \odot \dots \odot v^{*l_r} \otimes v^{*j} \otimes v_i,$$

with $\{v_1, \dots, v_n\}$ being the canonical basis of V and $\{v^{*1}, \dots, v^{*n}\}$ its dual basis.

Next lemma is an immediate consequence of the above remarks:

Lemma 4.3. *The mapping $(\pi_r^{r+1}, \Sigma^{(r+1)}) : J_0^{r+1}(FV) \rightarrow J_0^r(FV) \times V_1^{r+1,1}$ is a G -equivariant diffeomorphism.*

Now, let $\bar{\nabla}$ and ∇ be the canonical connections adapted to the $\{e\}$ -structure σ and the G -structure $s = q \circ \sigma$, respectively. The connection $\bar{\nabla}$ is characterized by the equations $\bar{\nabla}_{X_i} X_j = 0$, $1 \leq i, j \leq n$, so that the corresponding Christoffel symbols are given by $\bar{\Gamma}_{\alpha\beta}^\gamma = -\sum_i \frac{\partial \sigma_{\gamma i}}{\partial x^\alpha} \sigma^{i\beta}$. We define the tensor field F as the difference of both connections, i.e., by $F(X, Y) = \nabla_X Y - \bar{\nabla}_X Y$, $X, Y \in \mathfrak{X}(V)$, and denote by $F_{\alpha\beta}^\gamma = \Gamma_{\alpha\beta}^\gamma - \bar{\Gamma}_{\alpha\beta}^\gamma$, its components with respect to the canonical coordinates in V . From the expression in local coordinates of the connection ∇ given in Remark 2.1, it follows easily that these components have the form:

$$(10) \quad F_{\alpha\beta}^\gamma = \sum_{\lambda, \mu, \rho} \sigma^{\lambda\alpha} \sigma^{\mu\beta} \sigma_{\gamma\rho} (\sigma^* \omega)_\mu^\rho (X_\lambda) = \\ = \sum_{\lambda, \mu, \rho, i, j, k, h, \zeta} A_{\lambda\mu k}^{\rho ij} \sigma^{\lambda\alpha} \sigma^{\mu\beta} \sigma_{\gamma\rho} \sigma^{k\zeta} \left(\sigma_{hi} \frac{\partial \sigma_{\zeta j}}{\partial x^h} - \sigma_{hj} \frac{\partial \sigma_{\zeta i}}{\partial x^h} \right)$$

Due to the naturality of $\bar{\nabla}$ and ∇ , for each $g \in G$, the moving frame $R_g \circ (g^{-1} \cdot \sigma)$ has an associated tensor field $\bar{F}$ whose components are related to those of F by:

$$\bar{F}_{ij}^k(x) = g^{k\gamma} F_{\alpha\beta}^\gamma(g \cdot x) g_{\alpha i} g_{\beta j}.$$

Iterated derivation of these equations and evaluation at $x = 0$ yields:

$$\frac{\partial^k \bar{F}_{ij}^h}{\partial x^{p_1} \dots \partial x^{p_k}}(0) = g^{h\gamma} \frac{\partial^k F_{\alpha\beta}^\gamma}{\partial x^{l_1} \dots \partial x^{l_k}}(0) g_{l_1 p_1} \dots g_{l_k p_k} g_{\alpha i} g_{\beta j}.$$

So, we can also define an equivariant mapping $F^{(r)} : J_0^{r+1}(FV) \rightarrow V_1^{r,2}$ as follows:

$$F^{(r)}(j_0^{r+1} \sigma) = \sum_{l_1, \dots, l_r} \sum_{\alpha, \beta, \gamma} \frac{\partial^r F_{\alpha\beta}^\gamma}{\partial x^{l_1} \dots \partial x^{l_r}}(0) v^{*l_1} \odot \dots \odot v^{*l_r} \otimes v^{*\alpha} \otimes v^{*\beta} \otimes v_\gamma.$$

Lemma 4.4. *There exist a linear map $L^{(r)} : V_1^{r+1,1} \rightarrow V_1^{r,2}$ and a polynomial function $Q^{(r)} : J_0^r(FV) \rightarrow V_1^{r,2}$ in the standard coordinates of $J_0^r(FV)$ such that $F^{(r)} = L^{(r)} \circ \Sigma^{(r+1)} + Q^{(r)} \circ \pi_r^{r+1}$. Moreover, $L^{(r)}$ and $Q^{(r)}$ are G -equivariant.*

Proof. Derivation of (10) and evaluation at $x = 0$ show that the derivatives of $F_{\alpha\beta}^\gamma$ at $x = 0$ are of the form:

$$\frac{\partial^r F_{\alpha\beta}^\gamma}{\partial x^{l_1} \dots \partial x^{l_r}}(0) = \sum_{i,j,k} A_{\alpha\beta k}^{\gamma ij} \left(\frac{\partial^{r+1} \sigma_{kj}}{\partial x^{l_1} \dots \partial x^{l_r} \partial x^i}(0) - \frac{\partial^{r+1} \sigma_{ki}}{\partial x^{l_1} \dots \partial x^{l_r} \partial x^j}(0) \right) + O_r$$

where O_r stands for a polynomial in the derivatives up to order r of σ at $x = 0$.

Accordingly, we define $L^{(r)}$ as the linear map sending each element of $V_1^{r+1,1}$:

$$\sum_{l_1, \dots, l_{r+1}} \sum_{j,k} t_{l_1 \dots l_{r+1} j}^k v^{*l_1} \odot \dots \odot v^{*l_{r+1}} \otimes v^{*j} \otimes v_k,$$

to the element of $V_1^{r,2}$:

$$\sum_{l_1, \dots, l_r} \sum_{\alpha, \beta, \gamma} \sum_{i, j, k} A_{\alpha\beta k}^{\gamma ij} (t_{l_1 \dots l_r i j}^k - t_{l_1 \dots l_r j i}^k) v^{\star l_1} \odot \dots \odot v^{\star l_r} \otimes v^{\star \alpha} \otimes v^{\star \beta} \otimes v_{\gamma}.$$

Thus, $L^{(r)}$ is the composition:

$$V_1^{r+1,1} \xrightarrow{\delta^{r+1,1}} S^r(V^{\star}) \otimes \bigwedge^2 V^{\star} \otimes V^{1 \otimes (\delta^{-1} \circ P_{\text{im} \delta})} S^r(V^{\star}) \otimes V^{\star} \otimes \mathfrak{g} \subset V_1^{r,2},$$

where the first arrow stands for the Spencer's operator, which is defined in general as the map $\delta^{r+1, l-1} : S^{r+1}(V^{\star}) \otimes \bigwedge^{l-1} V^{\star} \otimes V \rightarrow S^r(V^{\star}) \otimes \bigwedge^l V^{\star} \otimes V$ given by:

$$(\delta^{r+1, l-1} t)(u_1, \dots, u_r, u'_1, \dots, u'_l) = \sum_{h=1}^l (-1)^{h+1} t(u_1, \dots, u_r, u'_h, u'_1, \dots, \widehat{u'_h}, \dots, u'_l),$$

for each $t \in S^{r+1}(V^{\star}) \otimes \bigwedge^{l-1} V^{\star} \otimes V$ and any vectors $u_1, \dots, u_r, u'_1, \dots, u'_l \in V$.

The difference $F^{(r)} - L^{(r)} \circ \Sigma^{(r+1)} : J_0^{r+1}(FV) \rightarrow V_1^{r,2}$ is a function depending polynomially in the derivatives up to order r of σ at 0, and hence there is a well-defined polynomial function $Q^{(r)} : J_0^r(FV) \rightarrow V_1^{r,2}$ such that $F^{(r)} - L^{(r)} \circ \Sigma^{(r+1)} = Q^{(r)} \circ \pi_r^{r+1}$.

Finally, it is clear that $L^{(r)}$ is G -equivariant, since it has been defined as a composition of homomorphisms of G -modules. The equivariance of the maps $F^{(r)}$, $L^{(r)}$, $\Sigma^{(r+1)}$ and π_r^{r+1} implies that $Q^{(r)}$ is also G -equivariant. ■

The desired characterization of the submanifold $\mathcal{S}^{r+1}$ is the following.

Theorem 4.5. *The submanifold $\mathcal{S}^{r+1} \subset J_0^{r+1}(FV)$ can be characterized as the set of $(r+1)$ -jets $j_0^{r+1}\sigma \in (\pi_r^{r+1}, \Sigma^{(r+1)})^{-1}(\mathcal{S}^r \times \bar{V}_1^{r+1,1})$ such that*

$$\text{sym}_1^{r,2} \left((L^{(r)} \circ \Sigma^{(r+1)}) (j_0^{r+1}\sigma) \right) = -\text{sym}_1^{r,2} \left((Q^{(r)} \circ \pi_r^{r+1}) (j_0^{r+1}\sigma) \right).$$

Proof. By definition, an r -jet $j_0^r\sigma \in J_0^r(FV)$ belongs to $\mathcal{S}^r$ if and only if $j_0^r\sigma = \Upsilon^r(j_0^r s)$ with $j_0^r s \in \mathcal{E}^r(V)_1$. Since Υ^r is a section of the projection $q^r : J_0^r(FV)_{u^0} \rightarrow J_0^r(FV/G)_{u^0}$, we have that $j_0^r s = q^r(j_0^r\sigma)$. So that $j_0^r\sigma \in \mathcal{S}^r$ if and only if $q^r(j_0^r\sigma) \in \mathcal{E}^r(V)_1$ and $(\Upsilon^r \circ q^r)(j_0^r\sigma) = j_0^r\sigma$.

Let us first study the second condition, which means that, up to order r , the moving frame σ is obtained by the parallel displacement of the canonical frame u^0 at $0 \in V$ using the canonical connection of the G -structure determined by σ . For each $j_0^r\sigma \in J_0^r(FV)_{u^0}$, let σ and s be local representatives of $j_0^r\sigma$ and $q^r(j_0^r\sigma)$, respectively, and let Γ_{ij}^k be the Christoffel symbols of the canonical connection $\nabla(s)$. Then, $(\Upsilon^r \circ q^r)(j_0^r\sigma) = j_0^r\sigma$ if and only if $j_0^r\sigma$ satisfies equation (9) for any $1 \leq k \leq r$, $1 \leq \gamma, i, i_1, \dots, i_r \leq n$.

Writing $\Gamma_{i_1\alpha}^\gamma = \bar{\Gamma}_{i_1\alpha}^\gamma + F_{i_1\alpha}^\gamma = -\sum_j \sigma^{j\alpha} \frac{\partial \sigma_{\gamma j}}{\partial x^{i_1}} + F_{i_1\alpha}^\gamma$, equation (9) reads:

$$\mathfrak{S}_{(i_1 \dots i_k)} \left(\frac{\partial^{k-1}}{\partial x^{i_2} \dots \partial x^{i_k}} \sum_{\alpha} F_{i_1\alpha}^\gamma \sigma_{\alpha i} \right) (0) = 0.$$

But this is equivalent to

$$\sum_{i_1, \dots, i_k} x^{i_1} \dots x^{i_k} \left(\frac{\partial^{k-1}}{\partial x^{i_2} \dots \partial x^{i_k}} \sum_{\alpha} F_{i_1\alpha}^\gamma \sigma_{\alpha i} \right) (0) = 0,$$

and hence to the equations

$$\left(\frac{d^{k-1}}{dt^{k-1}} \sum_{\alpha} \sigma_{\alpha i}(tx) \sum_{\beta} x^{\beta} F_{\beta \alpha}^{\gamma}(tx) \right) (0) = 0,$$

for any $1 \leq k \leq r$, $1 \leq \gamma, i, \leq n$ and any x in a neighborhood of $0 \in V$. Using induction, these equations are seen to be equivalent to the following:

$$\left(\frac{d^{k-1}}{dt^{k-1}} \sum_{\beta} x^{\beta} F_{\beta i}^{\gamma}(tx) \right) (0) = 0,$$

for any $1 \leq k \leq r$, $1 \leq \gamma, i, \leq n$ and any x , which can be written in terms of the partial derivatives at $0 \in V$ of F (and hence of σ) as:

$$(11) \quad \mathfrak{S}_{(i_1 \dots i_k)} \frac{\partial^{k-1} F_{i_1 \alpha}^{\gamma}}{\partial x^{i_2} \dots \partial x^{i_k}}(0) = 0,$$

for each $1 \leq k \leq r$, and for any indices $1 \leq \alpha, \gamma, i_1, \dots, i_k \leq n$.

Let us now find the equations provided by the condition $q^r(j_0^r \sigma) \in \mathcal{E}^r(V)_1$, which means that, up to order $r+1$, the normal coordinates of the connection attached to the G -structure $G \cdot \sigma$ are just the canonical coordinates of V . Setting $\Gamma_{i_1 i_2}^{\gamma} = \bar{\Gamma}_{i_1 i_2}^{\gamma} + F_{i_1 i_2}^{\gamma} = -\sum_j \sigma^{j i_2} \frac{\partial \sigma_{\gamma j}}{\partial x_1^i} + F_{i_1 i_2}^{\gamma} = \sum_j \sigma_{\gamma j} \frac{\partial \sigma^{j i_2}}{\partial x_1^i} + F_{i_1 i_2}^{\gamma}$ in (7) and using (11), we obtain that $q^r(j_0^r \sigma) \in \mathcal{E}^r(V)_1$ if and only if:

$$\left(\frac{d^{k-1}}{dt^{k-1}} \sum_j \sigma_{\gamma j}(tx) \frac{d}{dt} \left(\sum_{i_2} x^{i_2} \sigma^{j i_2}(tx) \right) \right) (0) = 0,$$

for any $1 \leq k \leq r$, $1 \leq \gamma \leq n$, and any x in a neighborhood of $0 \in V$. By induction, it follows again that these equations are equivalent to:

$$\left(\frac{d^k}{dt^k} \sum_{\beta} x^{\beta} \sigma^{\gamma \beta}(tx) \right) (0) = 0,$$

for any $1 \leq k \leq r$, $1 \leq \gamma \leq n$, and any x , which can be written as:

$$\mathfrak{S}_{(i_1 \dots i_{k+1})} \frac{\partial^k \sigma^{\gamma i_1}}{\partial x^{i_2} \dots \partial x^{i_{k+1}}}(0) = 0$$

or, equivalently, as:

$$(12) \quad \mathfrak{S}_{(i_1 \dots i_{k+1})} \frac{\partial^k \sigma_{\gamma i_1}}{\partial x^{i_2} \dots \partial x^{i_{k+1}}}(0) = 0$$

for each $1 \leq k \leq r$, and for any indices $1 \leq \gamma, i_1, \dots, i_{k+1} \leq n$. Finally, an r -jet $j_0^r \sigma \in J_0^r(FV)$ belongs to $\mathcal{S}^r$ if and only if it satisfies $\sigma_{\alpha \beta}(0) = \delta_{\alpha \beta}$, together with equations (12) and (11).

For the inductive definition of the submanifolds $\mathcal{S}^r$ appearing in the statement of the theorem, notice that an $(r+1)$ -jet $j_0^{r+1} \sigma \in J_0^{r+1}(FV)$ lies in $\mathcal{S}^{r+1}$ if and

only if $\pi_r^{r+1}(j_0^{r+1}\sigma) \in \mathcal{S}^r$ and equations (12) and (11) are satisfied at the top level, i.e.:

$$(13) \quad \mathfrak{S}_{(i_1 \dots i_{r+2})} \frac{\partial^{r+1} \sigma_{\alpha i_1}}{\partial x^{i_2} \dots \partial x^{i_{r+2}}}(0) = 0,$$

$$(14) \quad \mathfrak{S}_{(i_1 \dots i_{r+1})} \frac{\partial^r F_{i_1 \alpha}^\gamma}{\partial x^{i_2} \dots \partial x^{i_{r+1}}}(0) = 0,$$

for any indices $1 \leq \alpha, \gamma, i_1, \dots, i_{r+2} \leq n$.

Using the diffeomorphism given in Lemma 4.3 and the definition of the operators $F^{(r)}$, and taking into account that

$$\begin{aligned} & \text{sym}_q^{r,p} (v^{\star i_1} \odot \dots \odot v^{\star i_r} \otimes v^{\star i_{r+1}} \otimes \dots \otimes v^{\star i_{r+p}} \otimes v_{j_1} \otimes \dots \otimes v_{j_q}) \\ &= v^{\star i_1} \odot \dots \odot v^{\star i_r} \odot v^{\star i_{r+1}} \otimes \dots \otimes v^{\star i_{r+p}} \otimes v_{j_1} \otimes \dots \otimes v_{j_q} \\ &= \frac{1}{r+1} \mathfrak{S}_{(i_1 \dots i_{r+1})} v^{\star i_1} \odot \dots \odot v^{\star i_r} \otimes v^{\star i_{r+1}} \otimes \dots \otimes v^{\star i_{r+p}} \otimes v_{j_1} \otimes \dots \otimes v_{j_q}, \end{aligned}$$

we see that equations (13) and (14) are equivalent to $\text{sym}_1^{r+1,1}(\Sigma^{(r+1)}(j_0^{r+1}\sigma)) = 0$ and $\text{sym}_1^{r,2}(F^{(r)}(j_0^{r+1}\sigma)) = 0$, so each submanifold $\mathcal{S}^{r+1} \subset J_0^{r+1}(FV)$ is characterized by these conditions together with $\pi_r^{r+1}(j_0^{r+1}\sigma) \in \mathcal{S}^r$. That is:

$$\mathcal{S}^{r+1} = \left\{ j_0^{r+1}\sigma \in \left(\pi_r^{r+1}, \Sigma^{(r+1)} \right)^{-1} \left(\mathcal{S}^r \times \bar{V}_1^{r+1,1} \right) \mid \text{sym}_1^{r,2}(F^{(r)}(j_0^{r+1}\sigma)) = 0 \right\}$$

Finally, using the decomposition of $F^{(r)} = L^{(r)} \circ \Sigma^{(r+1)} + Q^{(r)} \circ \pi_r^{r+1}$ given in Lemma 4.4, we obtain the desired result. ■

5. DEFINITION OF THE G -MODULE STRUCTURE

In this section, we will define a G -module structure on each $\mathcal{S}^r$, inductively on r , such that the projections are homomorphisms of G -modules. The corresponding G -manifolds $\mathbf{S}^r = \mathcal{E}^r(V)/\mathfrak{G}_0^{r+1}$ will inherit G -module structures, and the natural projections between them will also be homomorphisms of G -modules. Therefore we will obtain a G -module structure on the projective limit $\mathbf{S}^\infty = \varprojlim \mathbf{S}^r$.

A main tool to define a G -module structure on each $\mathcal{S}^r$ (and therefore on each $\mathbf{S}^r$) is next lemma.

Lemma 5.1. *The projections $\pi_r^{r+1} : \mathcal{S}^{r+1} \rightarrow \mathcal{S}^r$ admit smooth G -equivariant sections $\mu_{r+1}^r : \mathcal{S}^r \rightarrow \mathcal{S}^{r+1}$.*

Proof. Let us denote by $\bar{L}^{(r)} : \bar{V}_1^{r+1,1} \rightarrow V_1^{r,2}$ the restriction of $L^{(r)}$ to $\bar{V}_1^{r+1,1}$, and $W^{r+1} := \ker(\text{sym}_1^{r,2} \circ \bar{L}^{(r)})$. Let us assume for the moment that we can choose a G -submodule $Z^{r+1} \subset \bar{V}_1^{r+1,1}$ such that $\bar{V}_1^{r+1,1} = W^{r+1} \oplus Z^{r+1}$. Then, the restriction $(\text{sym}_1^{r,2} \circ \bar{L}^{(r)})|_{Z^{r+1}} : Z^{r+1} \rightarrow \text{im}(\text{sym}_1^{r,2} \circ \bar{L}^{(r)})$ is an isomorphism of G -modules. Using that the projections $\pi_r^{r+1} : \mathcal{S}^{r+1} \rightarrow \mathcal{S}^r$ are onto, together with the fact that, by Theorem 4.5 we have that $\text{sym}_1^{r,2} \circ L^{(r)} \circ \Sigma^{(r+1)} = -\text{sym}_1^{r,2} \circ Q^{(r)} \circ \pi_r^{r+1}$ on $\mathcal{S}^{r+1}$ and $\Sigma^{(r+1)}(\mathcal{S}^{r+1}) \subset \bar{V}_1^{r+1,1}$, we obtain that

$$\begin{aligned} (-\text{sym}_1^{r,2} \circ Q^{(r)})(\mathcal{S}^r) &= (-\text{sym}_1^{r,2} \circ Q^{(r)} \circ \pi_r^{r+1})(\mathcal{S}^{r+1}) \\ &= (\text{sym}_1^{r,2} \circ L^{(r)} \circ \Sigma^{(r+1)})(\mathcal{S}^{r+1}) \subset \text{im}(\text{sym}_1^{r,2} \circ \bar{L}^{(r)}). \end{aligned}$$

Thus, for each $j_0^r \sigma \in \mathcal{S}^r$ there exists a unique $v^{r+1} \in Z^{r+1}$ such that

$$\left(\text{sym}_1^{r,2} \circ \bar{L}^{(r)} \right) (v^{r+1}) = -\text{sym}_1^{r,2} \left(Q^{(r)} (j_0^r \sigma) \right).$$

Now, using the isomorphism of Lemma 4.3, we set

$$\mu_{r+1}^r (j_0^r \sigma) = \left(\pi_r^{r+1} \times \Sigma^{(r+1)} \right)^{-1} (j_0^r \sigma, v^{r+1}).$$

It is clear that this defines a section $\mu_{r+1}^r : \mathcal{S}^r \rightarrow \mathcal{S}^{r+1}$ of the projection π_r^{r+1} . Moreover, from the G -equivariance of π_r^{r+1} , $\Sigma^{(r+1)}$, $\text{sym}_1^{r,2}$, $Q^{(r)}$, $L^{(r)}$ and the fact that Z^{r+1} is a G -submodule, it follows that μ_{r+1}^r is G -equivariant too.

It only remains to prove that it is indeed possible to choose a G -submodule $Z^{r+1} \subset \bar{V}_1^{r+1,1}$ such that $\bar{V}_1^{r+1,1} = W^{r+1} \oplus Z^{r+1}$. Note first that

$$(15) \quad V_1^{r+1,1} = \ker \left(\text{sym}_1^{r+1,1} \right) \oplus (S^{r+2} (V^*) \otimes V) = \bar{V}_1^{r+1,1} \oplus \ker \delta^{r+1,1}.$$

Therefore, $\delta|_{\bar{V}_1^{r+1,1}} : \bar{V}_1^{r+1,1} \rightarrow \text{im} \delta^{r+1,1}$ is an isomorphism of G -modules. It is easy to check that its inverse is given by $\left(\frac{r+1}{r+2} \right) \text{sym}_1^{r,2} : \text{im} \delta^{r+1,1} \rightarrow \bar{V}_1^{r+1,1}$.

Hence, it will enough to find a supplementary G -submodule $\hat{Z}^{r+1}$ of

$$\ker \left(\text{sym}_1^{r,2} \circ (1 \otimes \delta^{-1} \circ P_{\text{im} \delta}) \Big|_{\text{im} \delta^{r+1,1}} \right)$$

in $\text{im} \delta^{r+1,1}$. In order to do this, let us first prove that, if we consider $S^{r+1} (V^*) \otimes \mathfrak{g}$ as a G -submodule of $V_1^{r+1,1}$, then

$$\text{sym}_1^{r,2} \circ L^{(r)} \Big|_{S^{r+1} (V^*) \otimes \mathfrak{g}} = 1_{S^{r+1} (V^*) \otimes \mathfrak{g}}.$$

So let $\tau \in S^{r+1} (V^*) \otimes \mathfrak{g}$. Given vectors $u_1, \dots, u_{r+1}, u_{r+2} \in V$, we can consider $\tau(u_1, \dots, u_{r+1}) \in \mathfrak{g}$, and so $\tau(u_1, \dots, u_{r+1}) u_{r+2} \in V$. We have that

$$\begin{aligned} (\delta^{r+1,1} \tau) (u_1, \dots, u_{r+1}, u_{r+2}) &= \tau(u_1, \dots, u_{r+1}) u_{r+2} - \tau(u_1, \dots, u_{r+2}) u_{r+1} \\ &= \delta(\tau(u_1, \dots, u_r, \cdot))(u_{r+1}, u_{r+2}) \\ &= [(1_{S^r(V^*)} \otimes \delta) \tau] (u_1, \dots, u_r, u_{r+1}, u_{r+2}). \end{aligned}$$

Therefore, $\delta^{r+1,1} \tau = (1_{S^r(V^*)} \otimes \delta) \tau$, whence

$$(16) \quad \begin{aligned} \left(\text{sym}_1^{r,2} \circ L^{(r)} \right) \tau &= \left(\text{sym}_1^{r,2} \circ (1_{S^r(V^*)} \otimes \delta^{-1} \circ P_{\text{im} \delta}) \circ \delta^{r+1,1} \right) \tau \\ &= \text{sym}_1^{r,2} \tau = \tau. \end{aligned}$$

Thus,

$$\text{im} \left(\text{sym}_1^{r,2} \circ L^{(r)} \right) = S^{r+1} (V^*) \otimes \mathfrak{g}$$

and, according to (15), we also have that

$$\text{im} \left(\text{sym}_1^{r,2} \circ \bar{L}^{(r)} \right) = \text{im} \left(\text{sym}_1^{r,2} \circ L^{(r)} \right) = S^{r+1} (V^*) \otimes \mathfrak{g}.$$

Taking into account that, by (16), $\delta^{r+1,1}|_{S^{r+1}(V^*) \otimes \mathfrak{g}}$ is injective, we conclude that

$$\begin{aligned} &\dim \ker \left(\text{sym}_1^{r,2} \circ (1 \otimes \delta^{-1} \circ P_{\text{im} \delta}) \Big|_{\text{im} \delta^{r+1,1}} \right) \\ &= \dim (\text{im} \delta^{r+1,1}) - \dim (S^{r+1} (V^*) \otimes \mathfrak{g}) \\ &= \dim (\text{im} \delta^{r+1,1}) - \dim (\delta^{r+1,1} (S^{r+1} (V^*) \otimes \mathfrak{g})). \end{aligned}$$

Let us define $\hat{Z}^{r+1} = \delta^{r+1,1} (S^{r+1} (V^*) \otimes \mathfrak{g})$. Obviously it is a G -submodule of $\text{im} \delta^{r+1,1}$ of the required dimension. Moreover, (16) implies that

$$\hat{Z}^{r+1} \cap \ker \left(\text{sym}_1^{r,2} \circ (1 \otimes \delta^{-1} \circ P_{\text{im} \delta}) \Big|_{\text{im} \delta^{r+1,1}} \right) = \{0\}.$$

Hence, $\hat{Z}^{r+1}$ is the desired supplementary. ■

Proof of Theorem 1.1: The proof will be by induction on r . The case $r = 0$ is trivial: since S^0 has a unique element, determined by the conditions $\sigma_{ij}(0) = \delta_{ij}$, it admits the trivial G -module structure.

Now, assume that a G -module structure has been already defined on S^r and let us define one on S^{r+1} .

Let us consider the G -equivariant map $f^r = \Sigma^{(r+1)} \circ \mu_{r+1}^r : S^r \rightarrow \bar{V}_1^{r+1,1}$, and define a G -equivariant bijection $\psi^r : S^r \times \bar{V}_1^{r+1,1} \rightarrow S^r \times \bar{V}_1^{r+1,1}$ by $\psi^r(j_0^r \sigma, v^{r+1}) = (j_0^r \sigma, v^{r+1} - f^r(j_0^r \sigma))$, where the action of G on both sides is just the diagonal one.

After Lemma 5.1, we see that the image $(\pi_r^{r+1} \times \Sigma^{(r+1)})(S^{r+1})$ can be characterized as the set of pairs $(j_0^r \sigma, v^{r+1}) \in S^r \times \bar{V}_1^{r+1,1}$ satisfying:

$$\left(\text{sym}_1^{r,2} \circ \bar{L}^{(r)} \right) (v^{r+1}) = \left(\text{sym}_1^{r,2} \circ \bar{L}^{(r)} \right) (f^r(j_0^r \sigma))$$

(both members being equal to $-\text{sym}_1^{r,2}(Q^{(r)}(j_0^r \sigma))$). But this set is just the preimage by ψ^r of the G -submodule $S^r \oplus W^{r+1} \subset S^r \oplus \bar{V}_1^{r+1,1}$.

Now, we define the G -module structure on S^{r+1} as the one that makes linear the bijection:

$$(17) \quad \phi^r := \psi^r \circ \left(\pi_r^{r+1} \times \Sigma^{(r+1)} \right) : S^{r+1} \rightarrow S^r \oplus W^{r+1},$$

where we are considering the direct sum G -module structure on the right hand side.

The following diagram

$$\begin{array}{ccc} S^{r+1} & \xrightarrow{\phi^r} & S^r \oplus W^{r+1} \\ \pi_r^{r+1} \searrow & & \swarrow \text{pr}_1 \\ & S^r & \end{array}$$

where pr_1 denotes the projection onto the first summand, is commutative, so that the projections π_r^{r+1} are linear. Using induction, as well as the fact that ϕ^r is G -equivariant, we conclude that the action of G on each S^r is also linear. Moreover, the smoothness of the maps f^r implies the compatibility of the G -module structure with the manifold structure on each S^r . ■

Finally, the resulting G -module structure on each G -manifold S^r does not depend either on the supplementary subspace W used to define the canonical connections or on the sections μ_{r+1}^r between the corresponding spaces S^r . In fact, changing any of them would define isomorphic G -module structures, as follows from next general lemma:

Lemma 5.2. *Let M and N be two smooth n -dimensional manifolds endowed with smooth G -module structures. If M and N are diffeomorphic as G -manifolds, then they also are isomorphic as G -modules.*

Proof. Let $f : M \rightarrow N$ be a G -equivariant diffeomorphism. By composing, if necessary, with a translation in N we can assume that $f(0) = 0$. Due to the smoothness of the module structures, any linear isomorphisms $\varphi_1 : M \rightarrow \mathbb{R}^n$,

$\varphi_2 : N \rightarrow \mathbb{R}^n$ (onto $\mathbb{R}^n$ with its standard linear structure) define global charts of M and N . The map $\varphi_2 \circ f \circ \varphi_1^{-1} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a diffeomorphism leaving 0 fixed, and it is equivariant with respect to the linear actions of G on $\mathbb{R}^n$ induced by φ_1 and φ_2 . Then, the Fréchet derivative $D(\varphi_2 \circ f \circ \varphi_1^{-1})(0) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a linear G -equivariant isomorphism, and so is the composed map: $\varphi_2^{-1} \circ D(\varphi_2 \circ f \circ \varphi_1^{-1})(0) \circ \varphi_1 : M \rightarrow N$. ■

By definition, the G -module structure on $\mathcal{S}^{r+1}$ is isomorphic to $\mathcal{S}^r \oplus W^{r+1}$. This fact (together with the obvious identity $\mathcal{S}^0 = \{0\}$) yields an isomorphism of G -modules:

$$\mathbf{S}^r \cong W^1 \oplus W^2 \oplus \dots \oplus W^r,$$

and the G -module structure of $\mathbf{S}^r$ is determined by that of the spaces W^k , $k = 1, \dots, r$.

An element $t \in W^{r+1}$ lies in $\bar{V}_1^{r+1,1} := \ker(\text{sym}_1^{r+1,1})$, whence its components must satisfy the equations

$$(18) \quad \mathfrak{S}_{(i_1 \dots i_{r+1} i_{r+2})} t_{i_1 \dots i_{r+1} i_{r+2}}^k = 0, \quad 1 \leq k, i_1, \dots, i_{r+2} \leq n.$$

Denoting by $(L^{(r)}t)_{i_1 \dots i_{r+1} i_{r+2}}^k$ the components of $L^{(r)}t$, it follows that an element $t \in \bar{V}_1^{r+1,1}$ lies in W^{r+1} if and only if the following equations are also satisfied:

$$(19) \quad \mathfrak{S}_{(i_1 \dots i_{r+1})} (L^{(r)}t)_{i_1 \dots i_{r+1} i_{r+2}}^k = 0, \quad 1 \leq k, i_1, \dots, i_{r+2} \leq n.$$

Therefore, W^{r+1} is the G -submodule of $V_1^{r+1,1} = S^{r+1}(V^*) \otimes V^* \otimes V$ characterized by equations (18) and (19).

Notice that, in particular, $W^1 = \ker(\text{sym}_1^{0,2} \circ \bar{L}^{(0)})$. By definition, $\text{sym}_1^{0,2}$ is the identity in $V^* \otimes V^* \otimes V$. The mapping $\bar{L}^{(0)} : \bar{V}_1^{1,1} \rightarrow V_1^{0,2}$ is defined on $\bar{V}_1^{1,1} = \bigwedge^2 V^* \otimes V$ as

$$\bigwedge^2 V^* \otimes V \xrightarrow{P_{\text{im}\delta}} \delta(V^* \otimes \mathfrak{g}) \xrightarrow{\delta^{-1}} V^* \otimes \mathfrak{g}.$$

Thus $W^1 = \ker P_{\text{im}\delta} = W$.

Next, we will describe the G -modules W^r for some different choices of G .

Example 5.1 ($\{e\}$ -structures). For complete parallelisms, W^{r+1} is characterized by equation (18). Equation (19) is empty because the tensor F , and hence the operator $L^{(r)}$, are zero. Therefore, $W^{r+1} = \bar{V}_1^{r+1,1}$.

Example 5.2 ($O(n)$ -structures). We have seen (Example 2.1) that, for $G = O(n)$, the map δ is an isomorphism, with inverse $\delta^{-1} : \bigwedge^2 V^* \otimes V \rightarrow V^* \otimes \mathfrak{g}$ given, in components, by:

$$(\delta^{-1}(T))_{ij}^k = \frac{1}{2} (T_{ij}^k + T_{ki}^j + T_{kj}^i)$$

The operator $L^{(r)}$ is then given by:

$$\begin{aligned} (L^{(r)}t)_{i_1 \dots i_r i_{r+1} i_{r+2}}^k &= \frac{1}{2} \left(t_{i_1 \dots i_r i_{r+1} i_{r+2}}^k - t_{i_1 \dots i_r i_{r+2} i_{r+1}}^k \right. \\ &\quad + t_{i_1 \dots i_r k i_{r+1}}^{i_{r+2}} - t_{i_1 \dots i_r i_{r+1} k}^{i_{r+2}} \\ &\quad \left. + t_{i_1 \dots i_r k i_{r+2}}^{i_{r+1}} - t_{i_1 \dots i_r i_{r+2} k}^{i_{r+1}} \right). \end{aligned}$$

Taking the cyclic sum of this expression, and using (18), we obtain:

$$\begin{aligned} \mathfrak{S}_{(i_1 \dots i_{r+1})} (L^{(r)} t)_{i_1 \dots i_{r+1} i_{r+2}}^k &= \frac{1}{2} \left((r+1) t_{i_1 \dots i_r i_{r+1} i_{r+2}}^k + t_{i_1 \dots i_r i_{r+1} i_{r+2}}^k \right. \\ &\quad \left. - t_{i_1 \dots i_r i_{r+1} k}^{i_{r+2}} - (r+1) t_{i_1 \dots i_r i_{r+1} k}^{i_{r+2}} \right. \\ &\quad \left. + \mathfrak{S}_{(i_1 \dots i_{r+1})} \left(t_{i_1 \dots i_r k i_{r+2}}^{i_{r+1}} - t_{i_1 \dots i_r i_{r+2} k}^{i_{r+1}} \right) \right) \end{aligned}$$

Therefore, equations (19) can be written:

$$(20) \quad t_{i_1 \dots i_r i_{r+1} i_{r+2}}^k - t_{i_1 \dots i_r i_{r+1} k}^{i_{r+2}} = \frac{1}{r+2} \mathfrak{S}_{(i_1 \dots i_{r+1})} \left(t_{i_1 \dots i_r i_{r+2} k}^{i_{r+1}} - t_{i_1 \dots i_r k i_{r+2}}^{i_{r+1}} \right).$$

Taking the cyclic sum with respect to $i_1, \dots, i_{r+2}$ on both sides of last equation (using again (18)) yields:

$$\begin{aligned} - \mathfrak{S}_{(i_1 \dots i_{r+2})} t_{i_1 \dots i_r i_{r+1} k}^{i_{r+2}} &= \frac{1}{r+2} \left((r+1) \mathfrak{S}_{(i_1 \dots i_{r+2})} t_{i_1 \dots i_r i_{r+1} k}^{i_{r+2}} + \mathfrak{S}_{(i_1 \dots i_{r+2})} t_{i_1 \dots i_r i_{r+1} k}^{i_{r+2}} \right) \\ &= \mathfrak{S}_{(i_1 \dots i_{r+2})} t_{i_1 \dots i_r i_{r+1} k}^{i_{r+2}}, \end{aligned}$$

from which it follows that:

$$\mathfrak{S}_{(i_1 \dots i_{r+2})} t_{i_1 \dots i_r i_{r+1} k}^{i_{r+2}} = 0.$$

Last equation can be introduced in (20) giving:

$$t_{i_1 \dots i_r i_{r+1} i_{r+2}}^k - t_{i_1 \dots i_r i_{r+1} k}^{i_{r+2}} = \frac{1}{r+2} \left(-t_{i_1 \dots i_r i_{r+1} k}^{i_{r+2}} + t_{i_1 \dots i_r i_{r+1} i_{r+2}}^k \right),$$

from which one can conclude that the equations characterizing the subspace $W^{r+1} \subset V_1^{r+1,1}$ are (18) and

$$(21) \quad t_{i_1 \dots i_r i_{r+1} i_{r+2}}^k - t_{i_1 \dots i_r i_{r+1} k}^{i_{r+2}} = 0,$$

for any indices $1 \leq k, i_1, \dots, i_{r+2} \leq n$.

It should be pointed that, in [4], Epstein describes the space of ∞ -jets of Riemannian metrics g at a point, as follows. He writes the Taylor series for $g_{ij}(x)$ in normal coordinates: $\delta_{ij} + \sum_{r \geq 1} g_{ij i_1 \dots i_r} x^{i_1} \dots x^{i_r}$. The coefficients $g_{ij i_1 \dots i_r}$ satisfy the following conditions:

1. They are symmetric in the first two indices.
2. They are symmetric in the last r -indices.
3. $\mathfrak{S}_{(i_1 \dots i_{r+1})} g_{ii_1 \dots i_{r+1}} = 0$ for $1 \leq i, i_1, \dots, i_{r+1} \leq n$.

Then, he defines $f_r \in (V^*)^{\otimes(r+2)}$ by $f_r(v_i, v_j, v_{i_1}, \dots, v_{i_r}) = g_{ij i_1 \dots i_r}$, and proves that the set of $f_r \in (V^*)^{\otimes(r+2)}$ satisfying the symmetry conditions corresponding to 1-3 is an irreducible $GL(n, \mathbb{R})$ -module Y_r with Young diagram having r squares in the first row and 2 squares in the second one, except that if $r = 1$ then $Y_r = \{0\}$. Moreover, for any sequence $f_r \in Y_r$, $2 \leq r < \infty$, there is a Riemannian metric whose Taylor series gives the elements f_r , so that one can regard $\Pi_{r \geq 2} Y_r$ as the space of ∞ -jets of Riemannian metrics at a point.

The identification between Riemannian metrics g and sections s of the bundle $FV/O(n) \rightarrow V$ together with Lemma 5.2 yield an isomorphism of $O(n)$ -modules:

$\mathbf{S}^r \cong \Pi_{2 \leq k \leq r} Y_k$. Moreover, it is straightforward to check that the mapping $g_{jki_1 \dots i_{r+1}} \mapsto t_{i_1 \dots i_r i_{r+1} j}^k$ also defines an isomorphism of $O(n)$ -modules $W^{r+1} \cong Y_{r+1}$.

Example 5.3 ($O(p) \times O(q)$ -structures). From the expression of $\delta^{-1} \circ P_{\text{im}\delta}$ given in Example 2.2, it is immediate that the components of $L^{(r)}t$ are:

$$(22) \quad (L^{(r)}t)_{i_1 \dots i_r i_{r+1} i_{r+2}}^k = \frac{1}{2} \left(t_{i_1 \dots i_r i_{r+1} i_{r+2}}^k - t_{i_1 \dots i_r i_{r+2} i_{r+1}}^k + t_{i_1 \dots i_r k i_{r+1}}^{i_{r+2}} - t_{i_1 \dots i_r i_{r+1} k}^{i_{r+2}} + t_{i_1 \dots i_r k i_{r+2}}^{i_{r+1}} - t_{i_1 \dots i_r i_{r+2} k}^{i_{r+1}} \right)$$

if $i_{r+1}, i_{r+2}, k \in I_\alpha$, $\alpha = 1, 2$,

$$(23) \quad (L^{(r)}t)_{i_1 \dots i_r i_{r+1} i_{r+2}}^k = \frac{1}{2} \left(t_{i_1 \dots i_r i_{r+1} i_{r+2}}^k - t_{i_1 \dots i_r i_{r+2} i_{r+1}}^k - t_{i_1 \dots i_r i_{r+1} k}^{i_{r+2}} + t_{i_1 \dots i_r k i_{r+1}}^{i_{r+2}} \right)$$

if $i_{r+1} \in I_\alpha$, $i_{r+2}, k \in I_\beta$, $\alpha \neq \beta$, and

$$(L^{(r)}t)_{i_1 \dots i_r i_{r+1} i_{r+2}}^k = 0$$

if $i_{r+2} \in I_\alpha$, $k \in I_\beta$, $\alpha \neq \beta$.

As in the previous example, relation (22) leads to the equations (21) if $i_{r+1}, i_{r+2}, k \in I_\alpha$, $\alpha = 1, 2$. On the other hand, taking the cyclic sum with respect to $i_1, \dots, i_{r+1}$ in (23), and using (18), we obtain that equations (21) are also satisfied if $i_{r+1} \in I_\alpha$, $i_{r+2}, k \in I_\beta$, $\alpha \neq \beta$. In this way, the equations of $W^{r+1} \subset V_1^{r+1,1}$ turn out to be (18) and:

$$(24) \quad t_{i_1 \dots i_r i_{r+1} i_{r+2}}^k - t_{i_1 \dots i_r i_{r+1} k}^{i_{r+2}} = 0,$$

for any indices $i_1, \dots, i_{r+1} \in I_1 \cup I_2$, $i_{r+2}, k \in I_\alpha$, $\alpha = 1, 2$.

Example 5.4 ($\mathbb{R}^*$ -structures). The expression of $\delta^{-1} \circ P_{\text{im}\delta}$ given in Example 2.3 leads to the following expression of $L^{(r)}$:

$$(L^{(r)}t)_{i_1 \dots i_{r+1} i_{r+2}}^k = \frac{1}{n-1} \delta_{i_{r+2}}^k \sum_j \left(t_{i_1 \dots i_{r+1} j}^j - t_{i_1 \dots j i_{r+1}}^j \right).$$

Taking the cyclic sum with respect to $i_1, \dots, i_{r+1}$ (for $k = i_{r+2}$), and using (18), we obtain:

$$\begin{aligned} & \mathfrak{S}_{(i_1 \dots i_{r+1})} (L^{(r)}t)_{i_1 \dots i_{r+1} k}^k = \\ &= \frac{1}{n-1} \left((r+1) \sum_j t_{i_1 \dots i_r i_{r+1} j}^j + \sum_j t_{i_1 \dots i_r i_{r+1} j}^j \right) = \\ &= \frac{r+2}{n-1} \sum_j t_{i_1 \dots i_r i_{r+1} j}^j \end{aligned}$$

The resulting equations of W^{r+1} are then (18) and:

$$(25) \quad \sum_k t_{i_1 \dots i_{r+1} k}^k = 0, \quad 1 \leq k, i_1, \dots, i_{r+1} \leq n.$$

REFERENCES

- [1] I.M. Anderson, *Natural Variational Principles on Riemannian Manifolds*, Annals of Math. **120** (1984), 329–330.
- [2] D. Bernard, *Sur la Géométrie Différentielle des G -structures*, Ann. Inst. Fourier, Grenoble **10** (1960), 151–270.
- [3] S.S. Chern, *The Geometry of G -structures*, Bull. Amer. Math. Soc. **72** (1966), 167–219.
- [4] D.B.A. Epstein, *Natural Tensors on Riemannian Manifolds*, J. Differential Geom. **10** (1975), 631–645.
- [5] A. Ferrández and V. Miquel, *Hermitian Natural Tensors*, Math. Scand. **64** (1989), 233–250.
- [6] A. Fujimoto, *Theory of G -structures*, English Edition translated from the original Japanese, Publications of the Study Group of Geometry, vol. 1, 1972.
- [7] S. Kobayashi, *Transformation Groups in Differential Geometry*, Springer-Verlag, Berlin-Heidelberg-New York, 1972.
- [8] S. Kobayashi and K. Nomizu, *Foundations of Differential Geometry*, Volume I, John Wiley & Sons, New York, 1963.
- [9] T.Y. Thomas, *The Differential Invariants of Generalized Spaces*, Cambridge University Press, London, 1934.
- [10] A. Valdés, *Differential Invariants of $\mathbb{R}^*$ -Structures*, Math. Proc. Cambridge Philos. Soc. (1996), **119**, 341–356.

DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA, ESCUELA SUPERIOR DE INFORMÁTICA, UNIVERSIDAD COMPLUTENSE DE MADRID, 28040-MADRID, SPAIN
E-mail address: `celia_martinez@mat.ucm.es`

INSTITUTO DE FÍSICA APLICADA, CSIC, SERRANO 144, 28006-MADRID, SPAIN
E-mail address: `jaime@iec.csic.es`

DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA, FACULTAD DE CIENCIAS MATEMÁTICAS, UNIVERSIDAD COMPLUTENSE DE MADRID, 28040-MADRID, SPAIN
E-mail address: `avaldes@eucmos.sim.ucm.es`